





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 235/26, C07K 5/08 C07K 5/10, 5/06	A1	(11) International Publication Number: WO 91/01976 (43) International Publication Date: 21 February 1991 (21.02.91)
(21) International Application Number: PCT/CA90/00248 (22) International Filing Date: 3 August 1990 (03.08.90) (30) Priority data: 389,852 4 August 1989 (04.08.89) US (71) Applicant: IAF BIOCHEM INTERNATIONAL INC. [CA/CA]; 10900, rue Hamon, Montreal, Quebec H3M 3A2 (CA). (72) Inventors: BRILLON, Denis ; 119, de Galais, #2, Laval, Québec H7N 2Y6 (CA). SAUVE, Gilles ; 401, Sigmund- Freud, Laval, Quebec H7M 3X9 (CA). BOULOS, Zacha- rie ; 595, de l'Argentiére, #301, Laval, Quebec H7N 4A1 (CA). BELLEAU, Bernard (deceased).		(74) Agent: BROUILLETTE, Robert; Clark, Woods, 1245, Sherbrooke Street West, Suite 2000, Montreal, Quebec H3G 1G2 (CA). (81) Designated States: AT, AT (European patent), AU, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CM (OAPI patent), DE*, DE (European patent)*, DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), HU, IT (European patent), JP, KP, KR, LU, LU (European pa- tent), ML (OAPI patent), MR (OAPI patent), NL, NL (European patent), NO, RO, SD, SE, SE (European pa- tent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent). Published With international search report.

(54) Title: THIOACYLATING REAGENTS AND INTERMEDIATES, THIOPEPTIDES, AND METHODS FOR PREPARING AND USING SAME



(57) Abstract

Thioacylating reagents are provided for the introduction of thioamide bonds into growing peptides represented by structure (I), where the substituents are disclosed herein. Intermediate precursors for preparing these thioacylating reagents are also provided. A process for preparing the thioacylating reagents and the intermediate precursors is further provided. Thiopeptides, and salts thereof, that exhibit pharmacological utility are provided and are represented by formula (II), wherein R¹, R³, R⁴, X and n are as defined herein. Methods for preparing these thiopeptides from solution and solid phase syntheses, utilizing the thioacylating reagents disclosed herein, are also provided.

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Mónaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Fasso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LJ	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
DE	Germany, Federal Republic of	LU	Luxembourg	TC	Togo
DK	Denmark			US	United States of America

- 1 -

THIOACYLATING REAGENTS AND INTERMEDIATES,
THIOPEPTIDES, AND METHODS FOR
PREPARING AND USING SAME

5 Technical Field of the Invention

 This invention relates to novel α -amino acid derivatives used as thioacylating reagents, intermediates formed as precursors to these derivatives, thiopeptides of biological and medicinal
10 importance prepared through the use of these thioacylating reagents which introduce thioamide linkages into the growing peptide, and their methods of production. The thiopeptides disclosed herein are characterized by providing superior activity in vivo as
15 biological response modifiers, neuroeffectors, immunomodulators and the like and by increased resistance to enzymatic degradation due to the introduction of a thioamide linkage into the backbone of the peptide. The thiopeptides are described herein
20 as peptides which contain a thioamide linkage between adjacent amino acid residues. At least one such substitution by a sulphur atom of an oxygen atom in the amide bond of the backbone of the peptide structure is desired. As will be appreciated by the disclosure to
25 follow, the thioacylating reagents may be used to introduce this thioamide linkage into a growing peptide with relative ease and in substantially higher yields than observed in prior processes while retaining the

- 2 -

optical integrity of the peptide. The phrase growing peptide signifies that amino acid chain elongation is occurring which increases the size of the peptide by incorporating additional residues into the peptide
5 sequence.

Background of the Invention


The number of biologically active peptides is quite large. However, their potential utility as response modifiers, neuroeffectors or immunomodulators
10 is dramatically circumscribed by their demonstrated very short half-lives in vivo and their lack of effectiveness when administered orally. This latter phenomenon is primarily due to the extreme lability of biologically active polypeptides in the presence of the
15 peptidases and proteases normally found in the digestive tract.

It is desirable to stabilize the backbone amide linkages of these biologically active peptides against such proteolytic enzymes in order to improve
20 the pharmacokinetic properties of these peptides. Enhanced stability to enzymatic degradation would make these peptides more useful therapeutic agents.

Recent advances in chemical replacement or modification of peptide linkages indicate that such
25 linkage stabilization is feasible. By replacement of peptide linkages with thioamide bonds at those positions of the peptide backbone responsible for the biological response-limiting cleavage by peptidases and proteases, an increased stability to enzymatic
30 degradation is obtained for many thiopeptide analogs. Reid and von der Emden [W. Reid and W. von der Emden, "Aminosäure-thionester und Endothiopeptide, II", Liebigs Ann. Chem., 642, 128 (1961)] discusses racemic

- 3 -


thioamide formation through the thionester
thioacylating agent:



wherein R and R' are selected from lower alkyl and
aryl. Further, enhanced pharmacological activity is
10 exhibited for many of these analogues. Lajoie, et al.
(G. Lajoie, F. Lepine, S. LeMaire, F. Jolicoeur,
C. Aube, A. Turcotte and B. Belleau), "Synthesis and
Biological Activity of Monothionated Analogs of
Leucine-enkephalin", Int. J. Pept. Protein Res., 24,
15 316 (1984). Thiopeptide derivatives have demonstrated
increased activity in vivo as biological response
modifiers, neuroeffectors, and immunomodulators as
compared with their oxygenated analogs. For example,
Clausen, et al. [K. Clausen, A. Spatola, C. Lemieux,
20 P. Schiller, and S. Lawesson, "Evidence of a Peptide
Backbone Contribution Toward Selective Receptor
Recognition for Leucine Enkephalin Thioamide Analogs",
Biochem. Biophys. Res. Commun., 120, 305 (1984)]
demonstrates the increased pharmacological activity of
25 one such thiopeptide analog over its oxygenated
counterpart.

Methods for replacement for the carbonyl
oxygen atom of a carboxyl moiety with a sulphur atom
are known. Clausen, et al. [K. Clausen, M. Thorsen,
30 and S. Lawesson, "Studies on Amino Acids and Peptides.
Part 6. Methods for Introducing Thioamide Bonds into
the Peptide Backbone: Synthesis of the Four Monothio
Analogues of Leucine Enkephalin", J. Chem. Soc. Perkin
Trans. I, 785 (1984)] describes thioacylation by the
35 use of dithioesters of the formula:

- 4 -



5

wherein Z is carbobenzoxy and R is selected from hydrogen, lower alkyl, and aryl. No information, however, regarding racemization is described. It is also known that the thiopeptides so formed are useful reagents and intermediates for further thiopeptide synthesis. See, P. Campbell, and N. Nashed, "Carboxypeptidase A Catalyzed Hydrolysis of Thiopeptide and Thionester Analogues of Specific Substrates. An Effect on K_{cat} for Peptide, but not Ester, Substrates", J. Am. Chem. Soc., 104, 5221-26 (1982); P. Bartlett, K. Speer, and N. Jacobsen, "A Thioamide Substrate of Carboxypeptidase A", Biochemistry, 21, 1608-11, (1982); and L. Maziak, G. Lajoie, and B. Belleau, "Productive Conformation in the Bound State and Hydrolytic Behavior of Thiopeptide Analogues of Angiotensin-converting Enzyme Substrates", J. Am. Chem. Soc., 108, 182-83 (1986). Such thiopeptide derivatives also have shown resistance to enzymatic hydrolysis. W. Reid and E. Schmidt, "N-acylierte α -Aminoimidosäureester, Iminodipeptide und Endothiodipeptide", Liebigs Ann. Chem., 695, 217 (1966), for example, disclose the synthesis of a protected amino acid thionester as an intermediate in the preparation of a thiopeptide in moderate yield.

Thionation of peptides, or the replacement of an oxygen atom with a sulphur atom, at the carbonyl functionality of their peptide bonds has heretofore demonstrated a lack of reaction site specificity. Decreased overall yields have been observed because of side reactions and the by-products so formed which cause the purity of the product and the efficiency of

30

35

- 5 -

the reaction to suffer. Further, the optical integrity of the final product is often not maintained due to the reaction mechanism of the previously used thioacylating reagents. The limited effectiveness of these

5 thioacylating reagents severely circumscribed the potential of thiopeptides as pharmacological agents. Lack of an efficient method of producing pure, optically active thiopeptides has rendered the evaluation of pharmacological activity, stability to

10 enzymatic and pH degradation, and toxicity of such compounds very difficult, since sufficient quantities of these materials have heretofore been unobtainable.

The optical integrity of a compound relates to its ability to rotate light. This ability is

15 measured in an instrument known as a polarimeter which utilizes a zero point reference. The degree to which a chemically pure material rotates light indicates its relative optical purity. That is, a material may be chemically pure while being optically inactive or

20 racemic. The amount of activity that is observed from a material is often dependent upon its optical purity. Two enantiomers although possessing identical chemical formulae may have completely different biological activities. It is common in medicinal applications for

25 a compound of one optical configuration to exhibit activity and usefulness, while its optical rotamer or complementary enantiomer demonstrates a different activity or is wholly inert. Thus, where optical configuration is important, optical purity, as well as

30 chemical purity, is an important concern.

It is desirable that a thiopeptide meet several criteria to be suitable for pharmacological study. First, the thiopeptide should demonstrate an increased resistance to enzymatic degradation. Second,

35 the thiopeptide should elicit an enhanced biological

- 6 -

response over its oxygenated counterpart. Third, it must be safe for human ingestion. Fourth, the thiopeptide should be capable of being produced in quantities large enough to perform clinical studies.

5 Regarding the first three concerns, the characteristics described should be possessed as inherent properties of the thiopeptide which establish it as superior to other peptides not containing a thioamide linkage between adjacent amino acid residues.

10 With reference to the last criterion, it is advantageous to be capable of producing large quantities of material. Several factors are important with respect to this consideration. The process for producing the thiopeptide is preferably simple,
15 efficient and economical. That is, the reaction scheme of the process should contain few steps, afford high overall yields, and demonstrate minimal by-product formation. Moreover, the scheme should preferably utilize inexpensive reagents and materials. Further,
20 the method should ensure the optical integrity of the growing peptide by avoiding reactions that will racemize the compound. That is, a racemic mixture is likely not to fully exhibit the desired pharmacological response.

25 Prior thioacylation processes have suffered from being cumbersome and complicated. Moreover, they do not afford products with a high degree of optical integrity and provide inadequate overall yields of the thiopeptide.

30 Accordingly, there is a need for thioacylating reagents which permit the selective incorporation of thioamide linkages into growing peptides at specific residue linkages while utilizing efficient reaction conditions. There is also a need
35 for a thioacylation process which will retain the

- 7 -

optical integrity of the resulting peptide and will produce such peptide in high yields. There is additionally a need for methods for preparing thioacylating reagents capable of simple and economical
5 reaction with amino acids and peptides to produce thiopeptides. There is yet another need for thiopeptides, and methods to prepare them, having increased enzymatic stability and enhanced biological activity over their oxygenated analogs.


10

Summary of the Invention


It is an object of this invention to provide thioacylating reagents. It is also an object of this invention to provide thioacylating reagents which will introduce thioamide linkages into growing peptides in
15 high yield. It is a further object of this invention to provide thioacylating reagents which will retain the optical integrity of the peptide so formed. It is another object of this invention to provide novel intermediates to prepare the thioacylating reagents.
20 It is yet another object of this invention to provide thiopeptides which demonstrate greater pharmacological effectiveness with respect to activity and resistance to degradation than their oxygenated analogs. It is still another object of this invention to provide
25 methods for synthesizing these thioacylating agents, novel intermediates, and thiopeptides.

- 8 -

These and other objects are achieved herein
by thioacylating reagents represented by the formula:



wherein R^1 is hydrogen, C_1 - C_4 branched or unbranched
15 alkyl which may or may not be substituted by




wherein A is -O- or -NH-,
20 D is benzyl or xanthyl,
n is 1 or 2;

(b) $\begin{array}{c} \text{E} \\ | \\ -\text{CH}-\text{G}-\text{J}-\text{L} \end{array}$

25 wherein E is -H or - CH_3 ,
G is - CH_2 -, -O- or -S-,
J is -S- or - CH_2 -,
L is - CH_3 or phenyl;


- 9 -

(c)



wherein M and T may be the same or different and
 5 are selected from hydrogen, fluorine, chlorine, bromine
 and iodine and Q is hydrogen, hydroxy, or
 dichlorobenzoxy (2ClZ);


(d)



10

wherein V is carbobenzoxy or tosyl; or

(e)



15

R^2 is selected from t-butoxy, carbobenzoxy, chloro-
 benzyloxy, 9-fluorenylmethyloxycarbonyl, tosyl, trityl
 or xanthenyl;

20 R^3 is selected from hydrogen, methyl or ethyl; or
 R^1 and R^3 , taken together with the carbon atom to which
 they are attached, form a saturated hydrocarbon ring
 containing 3-5 carbon atoms;

R^4 is hydrogen; or R^1 and R^4 , taken together with the
 25 carbon and nitrogen atoms to which they are attached,
 form a saturated heterocycle containing 2-6 ring carbon
 atoms (i.e., an aziridine, azetidine, pyrrolidine, or
 piperidine ring); and

- 10 -

R⁵ is selected from hydrogen, fluorine, chlorine, bromine, iodine, amido, amino, carboxyl, carboxymethyl, cyano, guanido, hydroxyl, hydroxymethyl, mercapto, or nitro.

- 5 ~~_____~~ This invention also provides novel intermediate compounds that are extremely well suited for preparing the thioacylating reagent of this invention.

This invention additionally provides
10 thiopeptides that exhibit increased resistance to enzymatic degradation and demonstrate enhanced biological activity in vivo over their oxygen containing counterparts.

This invention further provides methods for
15 synthesizing these thioacylating reagents and novel intermediates useful in the preparation of these reagents, as well as processes for producing the thiopeptides represented herein.

In accordance with this invention, we provide
20 a class of reagents for thioacylation that may introduce thioamide linkages into peptides or other suitable compounds in a simple, efficient, economical manner and in high yield. Thioacylation carried out using these reagents will furthermore maintain the
25 optical integrity of the newly formed compound.


Also in accordance with this invention, we provide a method of producing these reagents and provide novel intermediates useful in the preparation of the thioacylating reagent.

30 In accordance with another aspect of this invention, we provide a series of thiopeptides which demonstrate an increased resistance to enzymatic degradation and an enhanced pharmacological activity as compared with corresponding peptides containing only
35 amide bonding between adjacent residues. These

- 11 -


thiopeptides show utility as biological response modifiers, neuroeffectors, immunomodulators and the like.

In accordance with this aspect of the invention, the thiopeptides are represented by the formula:



and salts thereof,

wherein R^1 is hydrogen, C_1 - C_4 branched or unbranched alkyl which may or may not be substituted by



wherein A is -O- or -NH-,

D is benzyl or xanthenyl,

20 n is 1 or 2;



wherein E is -H or - CH_3 ,


25 G is - CH_2 -, -O- or -S-,

J is -S- or - CH_2 -,

L is - CH_3 or phenyl;

(c)

30




- 12 -

wherein M and T may be the same or different and are selected from hydrogen, fluorine, chlorine, bromine and iodine and Q is hydrogen, hydroxy or 2ClZ;

5


(d)



wherein V is carbobenzoxy or tosyl; or

(e)

10




;

R³ is selected from hydrogen, methyl or ethyl;
R¹ and R³, taken together with the carbon atom to which they are attached, form a saturated hydrocarbon ring
15 containing 3-5 carbon atoms;
R⁴ is hydrogen or R¹ and R⁴, taken together with the carbon and nitrogen atoms to which they are attached, form a saturated heterocycle containing 2-6 ring carbon atoms (i.e., an aziridine, azetidine, pyrrolidine, or
20 piperidine ring); and n is 1-4.


- 13 -

Detailed Description of the InventionThioacylating Reagents

In accordance with the present invention, the thioacylating reagents are represented by:




wherein R^1 is represented by a member of the group consisting of hydrogen, C_1 - C_4 branched or unbranched alkyl which may or may not be substituted by a member selected from the group consisting of:



wherein A is -O- or -NH-,

D is benzyl or xanthenyl,

25 n is 1 or 2;



wherein E is -H or - CH_3 ,


30 G is - CH_2 -, -O- or -S-,

J is -S- or - CH_2 -,

L is - CH_3 or phenyl;


- 14 -

(c)



- 5 wherein M and T may be the same or different and are selected from hydrogen, fluorine, chlorine, bromine and iodine and Q is hydrogen, hydroxy or 2ClZ;


(d)



10

wherein V is carbobenzoxy or tosyl; or

(e)



15

R^2 is represented by a member selected from the class consisting of:

- t-butoxy, carbobenzoxy, chlorobenzyloxy, 9-fluorenylmethyloxycarbonyl, tosyl, trityl or xanthanyl;

20 R^3 is selected from hydrogen, methyl or ethyl; or R^1 and R^3 , taken together with the carbon atom to which they are attached, form a saturated hydrocarbon ring containing 3-5 carbon atoms;

- 25 R^4 is hydrogen; or R^1 and R^4 , taken together with the carbon and nitrogen atoms to which they are attached, form a saturated heterocycle containing 2-6 ring carbon


- 15 -

atoms (i.e., an aziridine, azetidine, pyrrolidine, or piperidine ring); and R⁵ is selected from hydrogen, fluorine, chlorine, bromine, iodine, amido, amino, carboxyl, carboxymethyl, cyano, guanido, hydroxyl, hydroxymethyl, mercapto, or nitro.

R¹ are typically substituents commonly found among natural α-amino acids. Particularly preferred groups include branched or unbranched alkyl groups which may or may not be substituted by amino, carboxy, guanido, hydroxy, hydroxymethyl or hydroxyphenyl.

The term thioacylating reagent is meant to include compounds which react with hydroxy and amino groups to introduce a thioacyl group to the nucleophilic substituent and becomes covalently bound thereto. The inclusion of thiocarbonyls in the amide bonds of peptides results in an increased resistance to hydrolysis and enzymatic destruction as compared with peptides of the same general structure but having conventional carbonyl moieties in their amide linkages.

The amino acid ortho amino thioanilides of this invention are represented by:



(II)


wherein R¹, R², R³, R⁴ and R⁵ in the compound are as defined hereinabove. Preferred values for R¹ in these thioanilide intermediates are substituents commonly found among natural α-amino acids. Particularly preferred groupings include branched or unbranched alkyl groups that may or may not be substituted by

- 16 -

amino, carboxy, guanido, hydroxy, hydroxymethyl or hydroxyphenyl.

The term amino-acid-ortho-amino-thioanilide is meant to include compounds having an ortho disubstituted amino benzene structure and an amino acid bound to said ortho disubstituted amino benzene structure by a thioamide linkage with said amino acid suitably protected at the amino terminus with an appropriate protecting group as defined for R^2 hereinabove.

The amino-acid-ortho-amino-anilides of this invention are represented by:




(I)

20 wherein R^1 , R^2 , R^3 , R^4 and R^5 in the compound are as defined hereinabove. Preferred values for R^1 are those substituents commonly found among natural α -amino acids. Particularly preferred groupings include branched or unbranched alkyl groups which may or may not be substituted by amino, carboxy, guanido, hydroxy, hydroxymethyl or hydroxyphenyl.


The term amino-acid-ortho-amino-anilide is meant to include compounds having an ortho disubstituted amino benzene structure and an amino acid bound to one of the amino substituents on said ortho disubstituted amino benzene structure through an amide linkage with said amino acid suitably protected at the amino terminus with an appropriate protecting group as defined for R^2 hereinabove.

- 17 -

The thioacylating reagents of the present invention may be prepared by initial reaction of an ortho-phenylenediamine conveniently with an amino acid according to the following scheme:



- 18 -



wherein R^1 , R^2 , R^3 , R^4 and R^5 are as defined hereinabove.

Ortho-phenylenediamine and amino acids

15 described herein may be reacted in the presence of a peptide coupling agent in a suitable solvent with stirring or agitation to form amino acid ortho amino anilides I. Surprisingly, however, selective amide formation occurs at only one of the two amino

20 substituents on the benzene ring. Contacting compounds of general formula I with a thionation reagent in the presence of a suitable solvent at -78°C to 0°C with stirring or agitation forms amino acid ortho amino thioanilides II. Subsequent treatment of compounds of

25 general formula II with a reagent well-suited to effect internal ring closure in a suitably inert solvent with stirring or agitation yields the desired compounds of general formula III. The reaction scheme to form the thioacylating reagents is illustrated hereinabove.

30 The process of peptide synthesis requires specific functional groups to react with other substituents to link amino acid residues in a desired manner to form a peptide with a sought after and known sequence. Since amino acids possess at least two

35 reactive substituents, the amine and carboxylic acid

- 19 -

portions, suitable protection or blocking of these functionalities is required to ensure that reaction will occur only at desired sites.

These protecting groups should be introduced to the moiety efficaciously while their removal should be performed under conditions which do not affect other portions of the molecule. In this manner, certain reactions and modifications may be performed on the amino acid, peptide, or other compound with assurance that the protected functionality will not interfere with the desired reaction. Further, by choosing a protecting group that is sensitive and labile to certain reactive conditions, a reaction scheme may be outlined to advantageously utilize these characteristics to effectively remove the protecting group once the synthesis is complete.

A variety of protecting groups known in the field of peptide synthesis and recognized by conventional abbreviations therein, may be found in T. Greene, Protective Groups In Organic Synthesis, Academic Press (1981). Among the preferred protecting groups that may be utilized for suitable protection of reactive nucleophilic substituents of R^1 are benzyl, carbobenzoxy or xanthenyl and for R^2 t-butoxy or carbobenzoxy.

Coupling of ortho-phenylenediamine with amino acids as described above to yield compounds of general formula I may be accomplished employing established techniques in the field of peptide chemistry. A broad range of suitable reactions are described in E. Gross & J. Meinhofer, 4 The Peptides: Analysis, Synthesis, Biology; Modern Techniques of Peptide and Amino Acid Analysis, John Wiley & Sons, (1981) and M. Bodanszky, Principles Of Peptide Synthesis, Springer-Verlag (1984). The peptide coupling agents which may be used

- 20 -

to assist condensation of amino and carboxylic acid moieties include N,N'-dicyclohexylcarbodiimide (DCC), N,N'-carbonyl diimidazole (CDI), 1-hydroxy benzotriazole (HOBt), ethyl chloroformate, and the like. A preferred technique uses DCC as the coupling reagent. The DCC method may be used with or without catalytic additives such as 4-dimethylaminopyridine (DMAP), copper (II) chloride or HOBt to hasten the reaction and suppress the racemization of the desired compound.

The DCC reaction is often performed at room temperature but may be performed from about -78°C to gentle reflux in a variety of solvents that are inert with respect to the reactants. The solvents are normally organic solvents which are polar and aprotic. Preferred solvents include dichloromethane, chloroform, diethyl ether, tetrahydroform (THF), N,N'-dimethylformamide (DMF), and the like. Particularly preferred solvents are dichloromethane and DMF. In general, the coupling reaction may be carried out at atmospheric pressure at a temperature of -78°C to reflux for a period of about 1-48h. Preferably, the reaction is carried out at -10°C to 25°C with stirring, shaking or agitation over a period of 4-6h.

Compounds of general formula II are typically prepared under anhydrous conditions, by reacting compounds of general formula I with a mixture of phosphorous pentasulfide and anhydrous sodium carbonate in an inert solvent. The reaction temperature is preferably about 0°C, but may then be varied from -78°C to gentle reflux. The solvent is preferably anhydrous THF, and other suitable solvents include dichloromethane, diethyl ether, DMF, and the like.

Compounds of general formula III may be prepared by contacting compounds of general formula II


- 21 -

with carbonyl ditriazole or phosgene in an inert solvent at a temperature of -78°C to gentle reflux, preferably room temperature. The solvent may be selected from, but is not limited to, dichloromethane, diethyl ether, DMF and THF.

In any of the synthesis methods described above, the desired products may be isolated from the reaction mixture by crystallization. Alternatively, chromatographic techniques including, but not limited to, normal phase, reverse-phase, ion-exchange, affinity, or gel permeation, may be employed, as well as electrophoresis or extraction or other means.


Thiopeptides

A novel class of thiopeptides contemplated by the present invention may be represented by the following formula:




(IV)

wherein X is S or O with at least one X being S, R^1 , R^3 , R^4 are as defined above, and n is 1-4. Preferred classes of thiopeptides are represented by the peptides comprising the following sequences:




(V)

- 22 -




wherein X, R^1 , R^3 , and R^4 are as defined above.

For compounds of formula V, the peptides may have four, three, or two thiocarbonyl moieties, but most preferably there will be one thiocarbonyl with the remaining values represented by X being oxygen. A particularly preferred thiopeptide of formula V for enhanced stability and increased pharmacological activity is represented by formula VII:



Similarly, for tetrapeptides of formula VI, especially preferred embodiments will be those wherein only one X represents a sulphur atom and the remaining two values for X each represent an oxygen atom. A most preferred series of thiopeptides of formula VI having increased resistance to enzymatic degradation and enhanced biological activity is represented by formula VIII:

- 23 -



(VIII)

The incorporation of thioamide linkages at positions on the peptide backbone which are susceptible to degradation may be performed in order to increase the peptide's resistance to enzymatic digestion. This enhanced stability may afford greater biological activities to the peptides prolonged existence. For example, thymopentin, which is merely a biologically active fragment of the polypeptide thymopoietin, may be modified to elicit this response. One such thymopentin derivative, 4-thiothymopentin, represented by formula VII, has exhibited approximately a three-fold increase in biological activity. A control group of nude athymic mice was administered 20 micrograms of thymopentin per subject according to well-known assay techniques. See, e.g., O. Archer, T. Pierce, B. Papermanter, and R. Good, "Reduced Antibody Response in Thymectomized Rabbits", *Nature*, 195, 191 (1962); D. Osoba and J. Miller, "Evidence for a Humeral Thymus Factor Responsible for the Maturation of Immunological Faculty", *Nature*, 199, 653 (1963); G. E. Ranges et al., "T-Cell Development in Normal and Thymopentin-treated Nude Mice", *J. Exp. Med.*, 156, 1057 (1982). The results of this assay showed a mean increase of 43% for the maturation of T cells and related immune responses. However, when the mice of the test group were

- 24 -

administered the same dosage under the same assay conditions, a mean increase of 128% for T cell maturation was observed. Thus, the increased effect in potency confirms that the introduction of a sulphur atom as a replacement for the carbonyl oxygen atom of an amide bond may decrease the rate of enzymatic degradation, enhance the affinity for relevant receptors, or act to assist both phenomena.

Other peptides that may be modified to incorporate thioamide linkages into the peptides backbone according to the thioacylation technology of the present invention include, but are not limited to, Vasopressin (9 amino acid residues), Somatostatin (14 amino acid residues), α -Melanotropin (13 amino acid residues), Leutienizing Hormone Releasing Hormone (LHRH, 10 amino acid residues), Adrenocorticotropin (ACTH, 39 amino acid residues), β -Endorphin (31 amino acid residues), and Atrial Natriuretic Factor (ANF, 33 amino acid residues).

Site specific and mild conditions which assist in the efficient introduction of a thioamide linkage to the backbone of a growing peptide chain may be accomplished in high yield by the use of 1-thioacyl-2-benzimidazolones as thioacylating reagents according to this invention. The resulting thiopeptides demonstrate increased stability and greater pharmacological potency while retaining the optical integrity of their component amino acid residues.

The thioacylation agents of the present invention permit the formation of thioamide linkages in growing peptides in substantially higher yields over those methods of thioamide introduction previously reported. The methods of the present invention yield thiopeptides that demonstrate previously unknown

- 25 -

stability with respect to resistance to enzymatic degradation. Further, the thiopeptides of this invention exhibit a substantial increase in pharmacological activity over peptide analogs that are
5 linked by amide bonding between residues.

According to this invention, a thioamide moiety may be introduced into a growing peptide at a specific site in the peptide sequence by reacting a thioacylating reagent according to the invention with
10 an amino acid or peptide. The peptide amino terminus must be protected at the terminal amino functionality. The reaction is advantageously carried out in a suitable solvent inert to the reactants in the presence of an appropriate peptide coupling reagent. The
15 preferred solvents include dichloromethane, chloroform, diethyl ether, THF, DMF, and the like. Particularly preferred solvents are dichloromethane and DMF. The preferred reaction conditions are from -78°C to gentle reflux for a period of about 1-48h. Particularly
20 preferred conditions are -10°C to 25°C stirring for a period of 4-6h.

Until the thioamide linkage is required to be introduced, the peptide may be synthesized under any peptide coupling conditions. Alternatively, the
25 thioamide linkage may be introduced first and the thiodipeptide so formed may then be enlarged employing generally recognized peptide coupling conditions.

Once the incorporation of the thioamide linkage is completed, and the thiopeptide is prepared,
30 the compound so formed may be entirely freed of its protecting groups according to well-known protocols such as treatment with liquid hydrogen fluoride (HF). Where, however, the peptide or thiopeptide formed requires selective removal of the protective groups,

- 26 -

usually from an amino terminus, suitable reaction conditions must be employed.

The t-butoxy (Boc) protected amino functionality of amino acid derivatives and terminal amines of peptides may be removed, e.g., by treatment with cold trifluoroacetic acid (TFA) at 0°C under suitable atmospheric conditions, e.g. adjusting the pH to about 8-9. The TFA salt of the amino acid derivative or protected peptide may be mixed in a suitable organic solvent and subjected to mild aqueous basic conditions. The organic solution, containing said amino acid derivative or said protected peptide with free amino functionality, may then be dried and concentrated to afford the free amino derivative.

In the case of 9-fluorenylmethyloxycarbonyl (Fmoc) protected amino acids, thiopeptides or Merrifield resin derivatives are presented, the corresponding free amino group may be generated selectively by treatment with piperidine in DMF under suitable atmospheric and thermal conditions.

An alternative synthetic approach for introducing thioamide linkages can be effectuated via the Merrifield solid phase methodology and its known variants. Thus, a Merrifield resin is prepared by well-known solid phase peptide synthesis methods. A covalently attached α -amino acid residue, attached at its carboxyl function or, similarly, a peptide with a free terminal amino functionality is carried by said resin. Treatment of said resin with said thioacylating reagent under standard solid phase peptide synthesis conditions affords the desired product.

Where this process introduces the thioamide linkage as the final step of thiopeptide formation, the thiopeptide may be liberated from said resin by using well-established methods. By employing liquid HF

- 27 -

containing dialkyl sulfide with anisole and thioanisole under suitable conditions at a temperature of -78°C to 0°C , the thiopeptide may be obtained free from all of the individual protecting groups of the component amino acid-residues.

The amounts of the reactants utilized in the aforementioned reaction may vary widely and the conditions required to facilitate reaction and encourage efficient completion may also vary widely. However, in general, the amounts of material employed to induce reaction in the processes discussed above will be substantially stoichiometric unless otherwise specified. In the following examples, reaction concentrations were generally held at 0.1 M to the reactants unless a higher concentration or dilution would be particularly useful for influencing the direction of a specific reaction. In practice, amounts will change depending upon variations in reaction conditions and the nature of the reactants.

The examples which follow are set forth to further illustrate various aspects of the present invention, but are not intended to limit its scope in any way.

EXAMPLE 1

25 Synthesis of 1-(α -N-Boc-L-seryl-O-benzyl-thioacyl)-2-benzimidazolone

a) Preparation of α -N-Boc-L-seryl-O-benzyl-ortho-amino-anilide

N-Boc-L-serine-O-benzyl ether (8.02 mmol) and ortho-phenylenediamine (11.6 mmol) were well dissolved in dichloromethane (21 ml) at 0°C and N,N'-dicyclohexylcarbodiimide (DCC) (8.27 mmol) added. The mixture was stirred for 1 hour at constant ice temperature and then filtered. The filtrate was

- 28 -

transferred to a separating funnel and washed successively with saturated brine/5% aqueous citric acid and saturated brine/5% aqueous sodium bicarbonate followed by saturated brine alone. The organic phase
5 was then dried, concentrated, and the residue purified by flash chromatography on silica gel employing a 3:1 hexane-ethyl acetate solvent as an eluant to yield the α -N-Boc-L-seryl-O-benzyl-ortho-amino-anilide as a solid in high yield (97%) which was then recrystallized to
10 analytical purity with a dichloromethane-pentane mixture. The compound was observed to have the following physical properties: Melting point 60-63°C; $[\alpha]_D^{20}$ (CHCl₃) -0.9; UV(CH₃CN) λ_{max} 293; Elemental composition (C₂₁H₂₇N₃O₄): Theoretical: C, 65.43; H, 7.06; N, 10.89. Found: C, 65.82; H, 7.42; N, 10.60.

Using the method of preparation described hereinabove and the appropriate starting materials, these additional ortho-amino anilides were synthesized:

- a1. α -N-Boc-L-alanyl-ortho-amino anilide
- 20 a2. α -N-Boc-L-arginyl-di-N-Cbz-ortho-amino anilide
- a3. α -N-Boc-L-arginyl-N-tosyl-ortho-amino anilide
- a4. α -N-Boc-L-asparaginy-N-xanthenyl-
25 ortho-amino anilide
- a5. α -N-Boc-L-aspartyl- β -benzyl ester-ortho-amino anilide
- a6. α -N-Fmoc-L-aspartyl- β -t-butyl ester-ortho-amino anilide
- 30 a7. α -N-Boc-L-cysteiny-N-benzyl ether-ortho-amino anilide
- a8. α -N-Boc-L-glutamyl- γ -benzyl ester-ortho-amino anilide
- a9. α -N-Boc-L-glutaminy-N-xanthenyl-ortho-
35 amino anilide

- 29 -

- a10. α -N-Boc-glycyl-ortho-amino anilide
 a11. α -N-Boc-L-histidyl-N-benzyl-ortho-amino
 anilide
 a12. α -N-Boc-L-histidyl-N-tosyl-ortho-amino
 5 anilide
 a13. α -N-Boc-L-isoleucyl-ortho-amino anilide
 a14. α -N-Boc-L-leucyl-ortho-amino anilide
 a15. α -N-Boc- ϵ -N-2ClZ-L-lysyl-ortho-amino
 anilide
 10 a16. α -N-Boc-L-methionyl-ortho-amino anilide
 a17. α -N-Boc-L-phenylalanyl-ortho-amino
 anilide
 a18. α -N-Boc-L-prolyl-ortho-amino anilide
 a19. α -N-Boc-L-threonyl-O-benzyl ether-
 15 ortho-amino anilide
 a20. α -N-Boc-L-tryptophyl-ortho-amino anilide
 a21. α -N-Boc-L-tyrosinyl-O-2,6-dichlorobenzyl
 ether-ortho-amino anilide
 a22. α -N-Boc-L-valyl-ortho-amino anilide
 20 Physical characterizations of these compounds
 are set forth in Table A.

b) Synthesis of α -N-Boc-L-seryl-O-benzyl-ortho-amino-thioanilide

- To freshly distilled tetrahydrofuran (THF)
 25 (67 ml) was added phosphorous pentasulfide (6.26 mmol)
 and anhydrous sodium carbonate (6.26 mmol). The
 mixture was permitted to stir at 20°C for 0.3 hours.
 The mixture was then cooled to 0°C followed by the
 addition of the N-Boc-L-seryl-O-benzyl-ortho-amino-
 30 anilide (of step a) (0.71 mmol). After standing at 0°C
 for 5-6 hours, 10% aqueous sodium phosphate (tribasic;
 22 ml) was added slowly followed by ethyl acetate
 (20 ml) and hexane (10 ml). The organic phase was
 separated, washed with brine, dried, and concentrated
 35 to yield an oil that was purified by flash

- 30 -

chromatography on silica gel using a 6:1:2 hexane-ethyl acetate-methylene chloride solvent mixture as the eluant to give the N-Boc-L-seryl-O-benzyl-ortho-amino thioanilide as a crystalline solid in a moderate yield
5 (50%). The compound was observed to have the following characteristics: Melting point 40-43°C; $[\alpha]_D^{20}$ (CHCl₃) -26.5; UV (CH₃CN) λ_{max} 271; Elemental composition (C₂₁H₂₇N₃O₄): Theoretical: C, 62.82; H, 6.80; N, 10.48; S, 7.98. Found: C, 63.06; H, 7.06; N, 10.42;
10 S, 8.23.

Using the method of preparation described hereinabove and the appropriate starting materials, these additional ortho amino thioanilides were synthesized:

- 15 b1. α -N-Boc-L-alanyl-ortho-amino thioanilide
- b2. α -N-Boc-L-arginyl-di-N-Cbz-ortho-amino-thioanilide
- b3. α -N-Boc-L-arginyl-N-tosyl-ortho-amino-
20 thioanilide
- b4. α -N-Boc-L-asparaginy-N-xanthenyl-ortho-amino-thioanilide
- b5. α -N-Boc-L-aspartyl- β -benzyl ester-ortho-amino-thioanilide
- 25 b6. α -N-Fmoc-L-aspartyl- β -t-butyl ester-ortho-amino-thioanilide
- b7. α -N-Boc-L-cysteiny-N-benzyl ether-ortho-amino-thioanilide
- b8. α -N-Boc-L-glutamyl- γ -benzyl ester-
30 ortho-amino-thioanilide
- b9. α -N-Boc-L-glutaminyl-N-xanthenyl-ortho-amino-thioanilide
- b10. α -N-Boc-glycyl-ortho-amino-thioanilide
- b11. α -N-Boc-L-histidyl-N-benzyl-ortho-amino-
35 thioanilide

- 31 -

b12. α -N-Boc-L-histidyl-N-tosyl-ortho-amino-thioanilide

b13. α -N-Boc-L-isoleucyl-ortho-amino-thioanilide

5 b14. α -N-Boc-L-leucyl-ortho-amino-thioanilide

b15. α -N-Boc- ϵ -N-2ClZ-L-lysyl-ortho-amino thioanilide

10 b16. α -N-Boc-L-methionyl-ortho-amino thioanilide

b17. α -N-Boc-L-phenylalanyl-ortho-amino-thioanilide

b18. α -N-Boc-L-prolyl-ortho-amino-thioanilide

15 b19. α -N-Boc-L-threonyl-O-benzyl-ether-ortho-amino thioanilide

b20. α -N-Boc-L-tryptophyl-ortho-amino thioanilide

20 b21. α -N-Boc-L-tyrosinyl-O-2,6-dichloro-benzyl-ether-ortho-amino-thioanilide

b22. α -N-Boc-L-valyl-ortho-amino thioanilide

Physical characterizations of these compounds are set forth in Table B.

25 c) Synthesis of 1-(α -N-Boc-L-seryl-O-benzyl-thioacyl)-2-benzimidazolone

The α -N-Boc-L-seryl-O-benzyl-ortho amino thioanilide (of step b) (3.11 mmol) and carbonyl ditriazole (4.36 mmol) were dissolved in THF (45 ml) and after stirring at 25°C for 6.5 hours, the solvent was removed in vacuo. The residue that remained was dissolved in dichloromethane (2 ml) and purified by flash chromatography. The product was eluted with 4:1 hexane-ethyl acetate to give pure 1-(N-Boc-L-seryl-O-benzyl-2-thioacyl)-2-benzimidazolone in high yield (91%). The compound was characterized by proton NMR

30
35

- 32 -

and the following physical characteristics were also recorded: Melting point 119-122°C; $[\alpha]_D^{20}$ (CHCl₃) -25.5; UV (CH₃CN) λ_{\max} 265; Elemental Composition (C₂₂H₂₅N₃O₄S): Theoretical: C, 61.81; H, 5.90; N, 9.82; S, 7.50. Found: C, 62.00; H, 6.01; N, 10.18; S, 7.30.

Using the method of preparation described hereinabove and the appropriate starting materials, these 1-(α -amino acid thioacyl)-2-benzimidazolone derivatives were synthesized:

- c1. 1-(α -Boc-L-alanyl-thioacyl)-2-benzimidazolone
- c2. 1-(α -Boc-L-arginyl-di-N-Cbz-thioacyl)-2-benzimidazolone
- 15 c3. 1-(α -Boc-L-arginyl-N-tosyl-thioacyl)-2-benzimidazolone
- c4. 1-(α -Boc-L-asparaginy-N-xanthenyl-thioacyl)-2-benzimidazolone
- c5. 1-(α -Boc-L-aspartyl- β -benzyl ester-thioacyl)-2-benzimidazolone
- 20 c6. 1-(α -Fmoc-L-aspartyl- β -t-butyl ester-thioacyl)-2-benzimidazolone
- c7. 1-(α -Boc-L-cysteiny-N-benzyl ether-thioacyl)-2-benzimidazolone
- 25 c8. 1-(α -Boc-L-glutamyl- γ -benzyl ester-thioacyl)-2-benzimidazolone
- c9. 1-(α -Boc-L-glutaminy-N-xanthenyl-thioacyl)-2-benzimidazolone
- c10. 1-(α -Boc-glycyl-thioacyl)-2-benzimidazolone
- 30 idazolone
- c11. 1-(α -Boc-L-histidyl-N-benzyl-thioacyl)-2-benzimidazolone
- c12. 1-(α -Boc-L-histidyl-N-tosyl-thioacyl)-2-benzimidazolone

- 33 -

- c13. 1-(α -Boc-L-isoleucyl-thioacyl)-2-benzimidazolone
- c14. 1-(α -Boc-L-leucyl-thioacyl)-2-benzimidazolone
- 5 c15. 1-(α -Boc-L-lysyl- ϵ -N-Cbz-thioacyl)-2-benzimidazolone
- c16. 1-(α -Boc-L-methionyl-thioacyl)-2-benzimidazolone
- c17. 1-(α -Boc-L-phenylalanyl-thioacyl)-2-benzimidazolone
- 10 c18. 1-(α -Boc-L-prolyl-thioacyl)-2-benzimidazolone
- c19. 1-(α -Boc-L-threonyl-O-benzyl ether-thioacyl)-2-benzimidazolone
- 15 c20. 1-(α -Boc-L-tryphophyl-thioacyl)-2-benzimidazolone
- c21. 1-(α -Boc-L-tyrosinyl-O-2,6-dichlorobenzyl ether-thioacyl)-2-benzimidazolone
- 20 c22. 1-(α -Boc-L-valyl-thioacyl)-2-benzimidazolone

Physical characterizations of these compounds are set forth in Table C.

EXAMPLE 2

Synthesis of 4-Thiothymopentin

25 a) Solution Synthesis

i) Preparation of α -N-Boc-L-valyl-L-tyrosyl-O-benzyl-benzyl ester-thioamide

α -N-Boc-L-tyrosyl-O-benzyl ether-benzyl ester
30 was treated with trifluoroacetic acid (TFA) at 0°C under nitrogen for 0.5 hours. The TFA was removed in vacuo to yield L-tyrosyl-O-benzyl-ether-benzyl ester TFA salt. This amino acid derivative was mixed in dichloromethane and treated with 5% aqueous sodium

- 34 -

bicarbonate. The organic phase was separated, dried, and concentrated to give the free amino derivative in quantitative yield.

L-tyrosyl-O-2,6-dichlorobenzyl ether benzyl ester (2 mmol) was dissolved in anhydrous N,N'-dimethylformamide (DMF) (0.5 ml) at 0°C under N₂ and 1-(α -N-Boc-valyl-thioacyl)-2-benzimidazolone (2.2 mmol) (from Example 1) was added in portions at 0°C with stirring over a 0.3 hour period. The mixture was stirred continuously at 0°C for 2 hours and allowed to warm to 25°C for 15-17 hours. The reaction was then filtered, concentrated in vacuo, the residue dissolved in ethyl acetate (15 ml) and the solution washed successively with 5% aqueous sodium bicarbonate, water, 5% aqueous citric acid, and water. The organic phase was then dried followed by evaporation and the residue placed on a flash column for purification. The protected dithiopeptide was eluted with a 3:2 ethyl acetate-hexane solvent mixture to afford α -N-Boc-L-valyl-L-tyrosyl-O-benzyl thioamide in good yield (80%). The compound was found to have the following physical characteristics: Melting point 56-58°C; IR (CHCl₃) 2972, 1735, 1500 cm⁻¹; UV (CHCl₃) λ_{max} 272.

ii) Preparation of α -N-Boc-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-3-thioamide

α -N-Boc-L-valyl-L-tyrosyl-O-benzyl thioamide (compound from (i)) was treated with TFA at 0°C for 0.5 hours under nitrogen to afford, after concentration in vacuo, L-valyl-L-tyrosyl-O-benzyl thioamide TFA salt. The compound was mixed in dichloromethane and treated with 5% aqueous sodium bicarbonate. The organic phase was separated, dried, and concentrated to give the free amino derivative in quantitative yield.

- 35 -

L-valyl-L-tyrosyl-O-benzyl-thioamide (2 mmol) was dissolved in anhydrous DMF (0.5 ml) at 0°C under N₂ and α -N-Boc-L-aspartyl- β -benzyl ester (2 mmol) was added to the solution with stirring. HOBt (2 mmol) and DCC (2 mmol) were added slowly at 0°C and stirring was allowed to continue overnight. The mixture was diluted with 8 volumes of ethyl acetate and the N,N'-dicyclohexylurea so formed was filtered away from the solution. The filtrate was transferred to a separatory funnel and washed successively with 5% aqueous sodium bicarbonate, 5% aqueous citric acid, and saturated brine. The organic phase was collected and dried over MgSO₄, filtered and concentrated in vacuo. The residue was purified by flash chromatography on silica gel employing a 1:1 hexane-ethyl acetate solvent mixture as an eluant to afford the α -N-Boc-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-3-thioamide in good yield (74%). The following physical characteristics were recorded for the compound: Melting point 112-114°C and UV (CHCl₃) λ_{max} 271.

iii) Preparation of α -N-Boc- ϵ -N-2ClZ-L-lysyl-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-L-3-thioamide

α -N-Boc-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-2-thioamide (compound from (ii)) was treated with TFA to remove the Boc group as in (ii) and L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-2-thioamide was obtained in quantitative yield.

L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-3-thioamide was dissolved in dry DMF (0.5 ml) at 0°C under N₂ and α -N-Boc- ϵ -2ClZ-L-lysine (2 mmol) was added to the solution with

- 36 -

stirring. HOBt (2 mmol) and DCC (2 mmol) were added slowly at 0°C and stirring was allowed to continue overnight. The mixture was diluted with 8 volumes of ethyl acetate and the N,N'-dicyclohexylurea so formed was filtered away from the solution. The filtrate was transferred to a separating funnel and washed successively with 5% aqueous sodium bicarbonate, 5% aqueous citric acid, and saturated brine. The organic phase was collected and dried over MgSO₄, filtered, and concentrated in vacuo. The residue was purified by flash chromatography on silica gel using a 1:1 hexane-ethyl acetate solvent mixture as an eluant to afford the α -N-Boc- ϵ -N-2ClZ-L-lysyl-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-3-thioamide in good yield (73%). The following physical characteristics were recorded for the compound: Melting point 71-73°C and UV (CHCl₃) λ_{max} 272.

iv) Preparation of α -N-Boc-N-tosyl-L-arginyl- ϵ -N-2ClZ-L-lysyl-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-4-thioamide

α -N-Boc- ϵ -N-2ClZ-L-lysyl-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-3-thioamide (compound from (iii)) was treated with TFA to remove the Boc group as in (ii) and ϵ -N-2ClZ-L-lysyl-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-3-thioamide was obtained in quantitative yield.

ϵ -N-2ClZ-L-lysyl-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-3-thioamide (2 mmol) was dissolved in dry DMF (0.5 ml) at 0°C under N₂ and α -N-Boc-N-tosyl-L-arginine (2 mmol) was added to the solution with stirring. HOBt (2 mmol) and DCC (2 mmol) were added slowly at 0°C and stirring was

- 37 -

allowed to continue overnight. The mixture was diluted with 8 volumes of ethyl acetate and the N,N'-dicyclohexylurea so formed was filtered away from the solution. The filtrate was transferred to a separating funnel and washed successively with 5% aqueous sodium bicarbonate, 5% aqueous citric acid, and saturated brine. The organic phase was collected and dried over MgSO_4 , filtered, and concentrated in vacuo. The residue was purification by flash chromatography using 9:1 chloroform-methanol as an eluant to afford the α -N-Boc-N-tosyl-L-arginyl- ϵ -N-2ClZ-L-lysyl-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-4-thioamide in good yield (78%). The following physical characteristics were recorded for the compound: Melting point 105-107°C; IR (CHCl_3) 1756, 1490 cm^{-1} ; UV (CHCl_3) λ_{max} 271. Elemental composition ($\text{C}_{71}\text{H}_{84}\text{Cl}_3\text{N}_9\text{O}_{14}\text{S}_2$): Theoretical: C, 58.13; H, 5.65; N, 8.34; S, 4.79. Found: C, 58.54; H, 5.81; N, 8.65; S, 4.40.

20 v) Preparation of 4-Thiothymopentin

α -N-Boc-N-tosyl-L-arginyl-L-lysyl- ϵ -N-2ClZ-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-benzyl ester-4-thioamide was dissolved in liquid hydrogen fluoride containing 10% by volume of anisole, ethyl methyl sulfide, and thioanisole. After 1 hour at 0°C, the solvent was evaporated in vacuo and the deprotected thiopeptide was purified by reversed phase chromatography on a Vydac C_{18} column utilizing 12% aqueous acetic acid as an eluant. The thiopeptide was characterized by proton NMR and was also found to possess the following physical properties: Melting point 146-148°C; UV (CHCl_3) λ_{max} 268; M/e 696.

- 38 -

Using the appropriate sequence of amino acid condensation reactions, the following mono-thiothymopentin analogs were prepared:

d1. L-arginyl-L-lysyl-L-aspartyl-L-valyl-L-tyrosine-1-thioamide

d2. L-arginyl-L-lysyl-L-aspartyl-L-valyl-L-tyrosine-2-thioamide

d3. L-arginyl-L-lysyl-L-aspartyl-L-valyl-L-tyrosine-3-thioamide

Physical characterizations for these thiopeptides are described in Table D.

b) Solid Phase Synthesis

i) Preparation of α -N-L-Boc-L-valyl-L-tyrosyl-O-benzyl-thioamide-resin ester

α -N-Boc-L-tyrosyl-O-benzyl ether attached to a benzyloxy group of a Merrified resin was treated with a 55% dichloromethane solution of TFA at room temperature for 1 hour. The resin was then collected, washed successively with four portions of 10 ml dichloromethane, four portions of 10 ml isopropanol (IPA) and dried for subsequent use.

L-tyrosyl-O-benzyl ether attached to a benzyloxy group of a Merrifield resin (0.632 mmol/g of resin) was added to a solution of 1-(α -N-Boc-valyl-thioacyl)-2-benzimidazolone (0.948 mmol) in dry DMF (7 ml) with stirring at 25°C. The reaction was stirred for 16 hours after which time another portion of the benzimidazolone (0.948 mmol) was added and stirring was resumed for 18 hours. The resin was collected, washed with four 10 ml portions of DMF, then four 10 ml portions of IPA and subsequently dried in preparation for further reaction.

- 39 -

ii) Preparation of α -N-L-Boc-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl-1-thioamide-resin ester

5 α -N-Boc-L-valyl-L-tyrosyl-O-benzyl ether-thioamide attached to a benzyloxy group of a Merrified resin was treated with a 55% dichloromethane solution of TFA at room temperature for 1 hour. The resin was then collected, washed successively with four portions
10 of 10 ml dichloromethane and four portions of 10 ml IPA and dried for subsequent use.

L-valyl-L-tyrosyl-O-benzyl ether-thioamide attached to a benzyloxy group of Merrifield resin (0.632 mmol/g of resin) was added to a solution of
15 α -N-Boc-L-aspartyl- β -benzyl ester (0.632 mmol) in DMF (7 ml) with stirring at 25°C. HOBt (0.632 mmol) and DCC (0.632 mmol) were added slowly with stirring. Reaction was allowed to proceed for 1-2 hours after which time the resin was collected, washed with four
20 10 ml portions of DMF, four 10 ml portions of IPA, and dried for further reaction.

iii) Preparation of α -N-Boc- ϵ -N-2ClZ-L-lysyl-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl-3-thioamide-resin ester

25

α -N-Boc-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-2-thioamide attached to a benzyloxy group of a Merrified resin was treated with a 55% dichloromethane solution of TFA at room temperature
30 for 1 hour. The resin was then collected, washed successively with four portions of 10 ml dichloromethane and four portions of 10 ml IPA and dried for subsequent use.

L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-2-thioamide attached to a benzyloxy
35

- 40 -

group of a Merrifield resin (0.632 mmol/g of resin) was added to a solution of α -N-Boc-N- ϵ -2ClZ-L-lysine (0.632 mmol) in DMF (7 ml) with stirring and the reaction proceeded for 1-2 hours. The resin was then collected, washed with four 10 ml portions of DMF, four 10 ml portions of IPA, and dried for further reaction.

iv) Preparation of α -N-Boc-N-tosyl-L-arginyl-L-lysyl- ϵ -N-2ClZ-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl-4-thioamide-resin ester

α -N-Boc- ϵ -N-2ClZ-L-lysyl-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-3-thioamide attached to a benzyloxy group of Merrified resin was treated with a 55% dichloromethane solution of TFA at room temperature for 1 hour. The resin was then collected, washed successively with four portions of 10 ml dichloromethane and four portions of 10 ml IPA and dried for subsequent use.

ϵ -N-2ClZ-L-lysyl-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-3-thioamide attached to a benzyloxy group of a Merrified resin (0.632 mmol/g of resin) was added to a solution of α -N-Boc-N-tosyl-L-arginine (0.632 mmol) in DMF (7 ml) with stirring at 25°C. HOBt (0.632 mmol) and DCC (0.632 mmol) were added slowly with stirring. Reaction was allowed to proceed for 1-2 hours. The resin was subsequently collected, washed with four 10 ml portions of IPA, and dried for final protecting group removal.

v) Preparation of 1-Thiothymopentin by Removal from Resin

α -N-Boc-N-tosyl-L-arginyl- ϵ -2ClZ-L-lysyl-L-aspartyl- β -benzyl ester-L-valyl-L-tyrosyl-O-benzyl ether-4-thioamide attached to a benzyloxy group of a

- 41 -

Merrifield resin (0.5 mmol) was treated with liquid hydrogen fluoride (5 ml) containing anisole, dimethyl sulfide, and thioanisole (0.5 ml 1:1:1 v/v) at 0°C for 1 hour. After evaporation of the solvent, the residue was dissolved in 10% aqueous acetic acid. The aqueous solution was washed with diethyl ether (30 ml), eluted with water and lyophilized to dryness. The crude thiopeptide was dissolved in 92% aqueous acetic acid (25 ml) and purified by reverse phase chromatography employing a C₁₈ packed column and the same acetic acid solvent as the eluant.

EXAMPLE 3

Preparation of 3-Thiotuftsia

a) Solution Synthesis

- 15 i) Synthesis of α -N-Boc-L-prolyl-N-tosyl-L-arginyl-benzyl ester thioamide

α -N-Boc-N-tosyl-L-arginyl-benzyl ester was treated with TFA at 0°C under nitrogen for 0.5 hours. The TFA was removed in vacuo to yield N-tosyl-L-arginyl-benzyl ester TFA salt. This amino acid derivative was mixed in dichloromethane and treated with 5% aqueous sodium bicarbonate. The organic phase was separate, dried, and concentrated to give the free amino derivative in quantitative yield.

N-tosyl-L-arginyl-benzyl ester (2 mmol) was dissolved in anhydrous DMF (0.5 ml) at 0°C under N₂ and 1-(α -N-Boc-L-thiopropyl)-2-benzimidazolone (2.2 mmol) (from Example 1) was added portionwise at 0°C with stirring over a 0.3 hour period. The mixture was stirred at 0°C continuously for 2 hours and permitted to warm to 25°C for 15-17 hours. The reaction was then filtered, concentrated in vacuo, the residue dissolved in ethyl acetate (15 ml) and the solution washed

- 42 -

successively with 5% aqueous sodium bicarbonate, water, 5% aqueous citric acid, and water. The organic phase was then dried followed by evaporation and the residue placed on a flash column for purification. The
5 protected thiodipeptide was eluted with a 3:2 ethyl acetate-hexane solvent mixture to afford α -N-Boc-L-thiopropyl-N-tosyl-L-arginyl-benzyl ester in high yield.

The recovered compound possessed the
10 following physical characteristics: Melting point 64-66°C and UV (CHCl_3) λ_{max} 270.

ii) Synthesis of α -N-Boc- ϵ -N-2ClZ-L-lysyl-L-prolyl-N-tosyl-L-arginyl-benzyl ester-2-thioamide

15 α -N-Boc-L-thiopropyl-N-tosyl-L-arginyl-benzyl ester was treated with TFA at 0°C under nitrogen for 0.5 hours. The TFA was removed in vacuo to yield L-thiopropyl-N-tosyl-L-arginyl-benzyl ester TFA salt. This thiodipeptide was mixed in dichloromethane and
20 treated with aqueous sodium bicarbonate. The organic phase was separated, dried, and concentrated to give the free amino derivative in quantitative yield.

L-thiopropyl-N-tosyl-L-arginyl-benzyl ester (2 mmol) was dissolved in anhydrous DMF (0.5 ml) at 0°C
25 under N_2 and α -N-Boc- ϵ -N-2ClZ-lysine (2 mmol) was added. HOBt (2 mmol) and DCC (2 mmol) were added slowly with stirring at 0°C and the reaction was allowed to continue overnight. The reaction was diluted with 8 volumes of ethyl acetate and the
30 N,N'-dicyclohexylurea so formed was filtered away from the mixture. The filtrate was transferred to a separatory funnel and washed successively with 5% aqueous sodium bicarbonate, 5% aqueous citric acid, and saturated brine. The organic phase was collected and
35 dried over MgSO_4 , filtered and concentrated in vacuo.

- 43 -

The residue was purified by flash chromatography on silica gel employing 1:1 hexane-ethyl acetate as an eluant to afford the α -N-Boc- ϵ -N-2ClZ-L-lysyl-L-prolyl-N-tosyl-L-arginyl-benzyl ester-2-thioamide in high yield.

The following physical characteristics were observed for the purified compound: Melting point 77-80°C; IR (CHCl₃) 1758, 1380 cm⁻¹; UV (CHCl₃) λ_{max} 271; Elemental composition (C₅₅H₇₁ClN₈O₁₁S₂):
Theoretical: C, 60.75; H, 7.10; N, 9.29. Found: C, 60.34; H, 7.07; N, 9.65.

iii) Synthesis of α -N-Boc-L-threonyl-O-benzyl ether- ϵ -N-2ClZ-L-lysyl-L-prolyl-N-tosyl-L-arginyl-benzyl ester-3-thioamide

α -N-Boc- ϵ -N-2ClZ-lysyl-L-prolyl-N-tosyl-L-arginyl-benzyl ester-2-thioamide was treated with TFA at 0°C under nitrogen for 0.5 hours. The TFA was removed in vacuo to yield ϵ -N-2ClZ-L-lysyl-L-prolyl-N-tosyl-L-arginyl-benzyl ester-2-thioamide TFA salt. This amino acid derivative was mixed in dichloromethane and treated with aqueous sodium bicarbonate. The organic phase was separated, dried, and concentrated to give the free amino derivative in quantitative yield.

ϵ -N-2ClZ-L-lysyl-L-prolyl-N-tosyl-L-arginyl-benzyl ester-2-thioamide (2 mmol) was dissolved in anhydrous DMF (0.5 ml) at 0°C under N₂ and α -N-Boc-threonine-O-benzyl ether (2 mmol) was added. HOBT (2 mmol) and DCC (2 mmol) were added portion-wise with stirring at 0°C and the reaction was allowed to continue overnight. The reaction was diluted with 8 volumes of ethyl acetate and the N,N'-dicyclohexylurea so formed was filtered away from the solution. The filtrate was transferred to a separatory funnel and

- 44 -

washed successively with 5% aqueous sodium bicarbonate, 5% aqueous citric acid, and saturated brine. The organic phase was collected and dried over MgSO_4 , filtered, and concentrated in vacuo. The residue was
5 purified by flash chromatography on silica gel employing a 1:1 hexane-ethyl acetate solvent mixture as an eluant to afford α -N-Boc-L-threonyl-O-benzyl ether- ϵ -N-2ClZ-L-lysyl-L-prolyl-N-tosyl-L-arginyl-benzyl ester-3-thioamide in high yield.

10 iv) Preparation of 3-Thiotuftsin

α -N-Boc-L-threonyl-O-benzyl ether- ϵ -N-2ClZ-L-lysyl-L-prolyl-N-tosyl-L-arginyl-benzyl ester-1-thioamide was dissolved in liquid hydrogen fluoride containing 10% by volume of anisole, ethyl methyl
15 sulfide, and thioanisole. After 1 hour at 0°C, the solvent was removed in vacuo and the deprotected thiopeptide was purified by reversed phase liquid column chromatography on a Vydac C_{18} column with 12% aqueous acetic acid as an eluant.

20 The thiopeptide exhibited the following physical characteristics: Melting point 169-171°C and UV (50% aqueous ethanol) λ_{max} 268.

The following mono-thiotuftsin analogs were prepared utilizing the appropriate starting materials
25 and the correct sequence of amino acid coupling reactions:

- e1. L-threonyl-L-lysyl-L-prolyl-L-arginine-2-thioamide
- e2. L-threonyl-L-lysyl-L-prolyl-L-arginine-
30 1-thioamide

Physical characterizations for these thiopeptides are described in Table E.

- 45 -

EXAMPLE 4Evaluation of Thymopentin and 4-Thiothymopentin
on T Cell Development in Nude Athymic Mice

This assay was performed according to the
5 protocol established by Lau and Goldstein. See C. Lau
and G. Goldstein, "Functional Effects of
Thymopoietin₃₂₋₃₆ (TP5) on Cytotoxic Lymphocyte
Precursor Units (CLP-U)", J. Immun., 124, 1861 (1980);
G. E. Ranges et al., "T Cell Development in Normal and
10 Thymopentin-treated Nude Mice", J. Exp. Med., 156, 1057
(1982).

The immunomodulatory action of thymopentin
and 4-thiothymopentin on immature T-cell development by
the expression of de novo antigens on T lymphocytes
15 after daily subcutaneous injections for two weeks in
four week old nude athymic mice was measured.
Following the final injection, the splenocytes were
prepared and radio-labelled to determine the percentage
of total cells bearing the radio labelled marker. Two
20 distinct sets of experiments established the
4-thiothymopentin as having a 128% and 227% increase of
cells bearing these markers over control animals
administered thymopentin at the same concentration and
rate, under the same conditions in this assay.

25 These findings suggest that thiothymopentin
analogs can induce a potent differentiation process
associated with an increase of cell surface markers of
T-lymphocytes.

Many variations and additional embodiments
30 will be readily apparent to those skilled in the art in
view of the foregoing disclosures and examples. For
example, thiopeptides of greater than five amino acid
residues having improved characteristics as described
herein, may be advantageously prepared according to the
35 methods of the present invention. All such obvious
variations and further embodiments are included within
the scope of the appended claims.

- 46 -

TABLE A

	Cmpnd	M.P. (°C)	$[\alpha]_D^{20}$ (CHCl ₃)	U.V. λ_{max} (CH ₃ CN)	Elemental Composition			
					%C %C	%H %H	%N %N	%S(found) %S(theor.)
5	a1	123-25	-54.5	294 (3.75)	60.23 60.20	7.77 7.58	14.76 15.03	
10	a2	116-18	-1.5	293 (3.48)	62.47 62.65	6.45 6.37	13.38 13.28	
	a3	101-02	-17.8	295 (3.60)	55.57	6.61	16.20	6.18
	a4	222-24	-10 (THF)	289 (3.58)	66.77 66.91	6.03 6.02	11.35 11.14	
15	a5	100-02	-13.0	292 (3.49)	64.16 63.90	7.05 6.58	10.56 10.16	
	a6	128-30	6.4	263 (4.13)	69.33 69.44	6.48 6.23	8.48 8.37	
20	a7	128-30	-14.2	293 (3.92)	63.00 62.82	7.01 6.78	10.77 10.46	7.54 7.98
	a8	75-78	-20.8	293 (3.39)	64.42 64.62	7.06 6.84	9.59 9.82	
	a9	212-13		294 (3.43)	67.59 67.43	6.54 6.24	11.14 10.84	
25	a10	148-49	0	293 (3.39)	58.56 58.86	7.31 7.22	15.44 15.83	
	a11	49-52	-5.2	294 (3.44)	65.90 66.19	7.11 6.71	15.65 16.07	

- 47 -

TABLE A (cont'd)

	Cmpnd	M.P. (°C)	$[\alpha]_D^{20}$ (CHCl ₃)	U.V. λ_{max} (CH ₃ CN)	Elemental Composition			
					%C %C	%H %H	%N %N	%S(found) %S(theor.)
5	a12	82-84	7.8	292 (3.45)				
10	a13	150-51	-38.8	294 (3.55)	63.50 63.53	8.54 8.47	13.28 13.08	
	a14	225-27	-2.5 (THF)	280 (3.07)	63.49 63.53	8.79 8.47	13.02 13.08	
	a15	94-97	-26.8	293 (3.63)	59.99 59.46	6.94 6.58	10.80 11.09	
15	a16	137-38	-29.0	294 (3.15)	56.49 56.61	7.69 7.42	12.11 12.39	9.34 9.44
	a17	141-42	0	301 (3.60)	67.38 67.58	7.42 7.09	11.88 11.81	
20	a18	164-66	-105	293 (3.52)	63.31 62.93	7.90 7.59	13.61 13.75	
	a19	142-44	7.2	292 (3.54)	65.95 66.14	7.52 7.32	10.37 10.51	
	a20	149-52	-2.6 (CH ₃ CN)	288 (4.17)	67.29 66.98	7.05 6.64	14.37 14.20	
25	a21	173-75	20.9 (THF)	293 (3.56)	60.70 61.14	5.67 5.51	7.62 7.92	
	a22	125-26	-39.8	292 (3.49)	62.94 62.54	8.32 8.19	13.81 13.68	

- 48 -

TABLE B

	Compnd	M.P. (°C)	$[\alpha]_D^{20}$ (CHCl ₃)	U.V. λ_{max} (CH ₃ CN)	Elemental Composition			
					%C %C	%H %H	%N %N	%S(found) %S(theor.)
5	b1	126-28	-71.4	270 (3.96)	57.14 56.92	7.27 7.17	14.48 14.22	10.96 10.85
10	b2	68-70	-14.6	271 (4.06)	61.08 61.09	6.48 6.21	12.92 12.95	4.70 4.94
	b3	130-37	-11.3	271 (4.23)	53.66 53.91	6.80 6.41	15.51 15.71	12.33 12.00
	b4	98-99	18.9	278 (4.12)	64.81 64.84	6.05 5.83	11.14 10.80	6.19 6.18
15	b5	125-26	-27.9	274 (4.16)	61.12 61.52	6.19 6.29	9.39 9.78	7.34 7.46
	b6	74-78	-12.8	264 (4.21)	67.13 67.29	6.47 6.04	8.45 8.12	5.92 6.19
20	b7	114-15	-48.0	273 (4.06)	60.30 60.40	6.58 6.52	9.86 10.06	15.02 15.35
	b8	51-53	-19.4	272 (3.75)	63.08 62.58	6.80 6.59	9.39 9.47	6.96 7.23
	b9	210-11 (THF)	-4.5	277 (4.20)	65.16 65.39	6.45 6.05	10.85 10.51	6.01 6.02
25	b10	124-25	-0.1	296 (4.00)	55.89 55.49	7.10 6.81	14.80 14.92	11.57 11.39
	b11	60-62	-21.0	273 (3.93)	65.04 63.83	6.82 6.47	15.63 15.50	7.04 7.10

- 49 -

TABLE B (cont'd)

	Compnd	M.P. (°C)	$[\alpha]_D^{20}$ (CHCl ₃)	U.V. λ_{max} (CH ₃ CN)	Elemental Composition			
					%C %C	%H %H	%N %N	%S(found) %S(theor.)
5	b12	116-17	81.2	268 (3.92)				
10	b13	139-40	-6.8	273 (4.08)	60.85 60.52	8.24 8.06	12.75 12.44	9.28 9.50
	b14	66-70	-42.9	273 (3.93)	60.65 60.52	8.25 8.06	12.51 12.44	9.27 9.50
	b15	56-58	-34.0	272 (3.98)	58.00 57.62	6.67 6.37	11.10 10.74	6.27 6.15
15	b16	46-48	-6.1	273 (3.98)	53.80 54.06	6.81 7.09	11.86 11.82	17.79 18.04
	b17	66-69	41.3	273 (4.06)	65.03 64.70	7.00 6.78	16.09 11.30	8.41 8.61
20	b18	73-75	-179 (c=0.5)	270 (4.10)	60.00 59.80	7.65 7.21	13.00 13.06	9.60 9.97
	b19	49-52	-34.6	272 (3.93)	63.74 63.52	7.57 7.33	9.90 10.10	7.46 7.71
	b20	161-62	19.9	273 (4.31)	64.52 64.36	6.59 6.38	13.76 13.64	7.63 7.81
25	b21	164-65	39.2	273 (4.14)	59.68 59.33	5.66 5.35	7.74 7.68	5.74 5.87
	b22	117-19	-7.7	273 (4.18)	59.63 59.40	7.72 7.79	12.74 12.98	10.03 9.91

- 50 -

TABLE C

	Compnd	M.P. (°C)	$[\alpha]_D^{20}$ (CHCl ₃)	U.V. λ_{max} (CH ₃ CN)	Elemental Composition			
					%C %C	%H %H	%N %N	%S(found) %S(theor.)
5	c1	103-05	-19.0	263 (3.60)	55.82 56.06	5.97 5.96	12.73 13.07	10.04 9.98
10	c2	152-54	3.4	263 (4.03)	60.35 60.52	5.80 5.68	12.41 12.45	4.62 4.75
	c3			NOT ISOLATED				
	c4	177-81	55.2 (DMF)	264 (3.74)				
	c5			NOT ISOLATED				
15	c6			NOT ISOLATED				
	c7	136-38	46.0	266 (3.78)	59.20 59.57	5.80 5.68	9.41 9.47	14.40 14.45
	c8	124-27	48.0	264 (3.92)	61.66 61.39	6.05 5.80	8.53 8.55	7.00 6.83
20	c9	132-34	53.5 (THF)	264 (3.79)	64.54 64.80	5.80 5.41	9.83 10.02	5.96 5.74
	c10	121-24	0	263 (3.92)	54.91 54.71	5.84 5.57	13.17 13.66	10.03 10.42
	c11			NOT ISOLATED				
25	c12			NOT ISOLATED				
	c13	72-73	129	265 (3.92)	59.79 59.48	7.18 6.93	11.86 11.55	8.87 8.82

- 51 -

TABLE C (cont'd)

5	Compnd	M.P. (°C)	$[\alpha]_D^{20}$ (CHCl ₃)	U.V. λ_{max} (CH ₃ CN)	Elemental Composition			
					%C %C	%H %H	%N %N	%S(found) %S(theor.)
10	c14	123-16	46.8	264 (3.90)	59.70 59.48	7.09 6.93	11.75 11.55	9.12 8.82
	c15	134-36	30.3	264 (4.00)	56.72 57.08	5.83 5.71	10.16 10.24	5.53 5.86
	c16	118-20	46.9	263 (4.10)	53.38 53.52	6.39 6.07	11.50 11.10	16.74 16.81
15	c17	166-69	141.3	263 (4.27)	63.32 63.45	5.73 6.83	10.87 10.46	8.26 8.06
	c18	136-38	-203	266 (3.94)	58.48 58.76	6.34 6.10	12.14 12.09	9.47 9.22
	c19	138-40	43.3	266 (4.01)	62.22 62.56	6.40 6.16	9.27 9.51	7.54 7.26
20	c20	164-67	146	266 (4.08)	63.55 63.28	5.81 5.54	13.00 12.83	7.29 7.34
	c21	187-89	109	264 (4.08)	58.66 58.74	4.99 4.75	7.76 7.34	5.47 5.60
	c22	148-50	146	266 (3.98)	58.79 58.42	6.94 6.63	11.86 12.02	9.11 9.17

- 52 -

TABLE D

	<u>Cmpnd</u>	<u>M.P.</u> <u>(°C)</u>	<u>U.V.</u> <u>λ_{max}</u> <u>(CH₃CN)</u>
5	d1	140-42	268 (3.86)
	d2	133-34	269 (3.71)
10	d3	148-50	269 (3.89)


TABLE E

	<u>Cmpnd</u>	<u>M.P.</u> <u>(°C)</u>	<u>U.V.</u> <u>λ_{max}</u> <u>(CH₃CN)</u>
15	e1	182-83	267 (3.81) (in 50% EtOH)


- 53 -

WHAT IS CLAIMED IS:

1. A thioacylating reagent of the formula:




wherein R^1 is hydrogen or C_1 - C_4 branched or unbranched alkyl which may be substituted by a substituent selected from the group consisting of:



wherein A is -O- or -NH-,

D is benzyl or xanthenyl,

n is 1 or 2;




wherein E is -H or -CH₃,

G is -CH₂-, -O- or -S-,


J is -S- or -CH₂-

L is -CH₃ or phenyl;




- 54 -

wherein M and T may be the same or different and are members selected from the group consisting of hydrogen, fluorine, chlorine, bromine and iodine, and Q is hydrogen, hydroxy, or dichlorobenzyloxy,



wherein V is carbobenzyloxy or tosyl, or



;

R^2 is selected from the group consisting of t-butoxy, carbobenzyloxy, chlorobenzyloxy, 9-fluorenylmethoxycarbonyl, tosyl, trityl and xanthenyl;

R^3 is hydrogen, methyl, or ethyl; or R^1 and R^3 , taken together with the carbon atom to which they are attached, form a cycloalkyl group containing 3-5 carbon atoms;

R^4 is hydrogen; or R^1 and R^4 , taken together with the carbon and nitrogen atoms to which they are attached, form a saturated heterocycle containing not less than 2, but not more than 6 ring carbon atoms; and


R^5 is selected from the group consisting of hydrogen, methyl, ethyl, fluorine, chlorine, bromine, iodine, amino, amido, azido, hydroxy, hydroxymethyl, carboxy, carboxymethyl, cyano, guanido, mercapto, and nitro.

- 55 -

2. The thioacylating reagent according to claim 1, wherein R^1 is C_1-C_4 branched or unbranched alkyl, R^2 is t-butoxy, and R^3 is hydrogen.

3. The thioacylating reagent according to claim 2, wherein R^1 and R^4 , taken together with the carbon and nitrogen atoms to which they are attached, form a saturated heterocycle containing 4 ring carbon atoms (a pyrrolidine ring), R^2 is t-butoxy, and R^3 and R^5 are hydrogen.

4. The thioacylating reagent according to claim 1, wherein R^1 is represented by the formula:



wherein A is -O- or -NH-,


D is benzyl or xanthenyl,

n is 1 or 2;

R^2 is t-butoxy; and

R^3 and R^5 are hydrogen.

5. A compound according to claim 1, wherein R^1 is represented by the formula:



wherein E is -H or -CH₃,

G is -CH₂-, -O- or -S-,

J is -S- or -CH₂-,


L is -CH₃ or phenyl;

R^2 is t-butoxy; and

R^3 and R^5 are hydrogen.

- 56 -

6. The thioacylating reagent according to claim 1, wherein R^1 is represented by the formula:




wherein M and T may be the same or different and are substituents selected from the group consisting of hydrogen, fluorine, chlorine, bromine and iodine, and Q is hydrogen, hydroxy, or $2ClZ$;

R^2 is t-butoxy; and

R^3 and R^5 are hydrogen.

7. The thioacylating reagent according to claim 1, wherein R^1 is represented by the formula:




wherein V is carbobenzyloxy or tosyl;

R^2 is t-butoxy; and

R^3 and R^5 are hydrogen.


8. The thioacylating reagent according to claim 1, wherein R^1 is:



- 57 -


R^2 is t-butoxy; and
 R^3 and R^5 are hydrogen.

9. The thioacylating reagent according to claim 1, wherein R^1 is represented by the formula:



wherein n is 1 - 4 and R^3 and R^5 are hydrogen.

10. The thioacylating reagent according to claim 1, wherein R^1 is:



R^2 is 9-fluorenylmethyloxycarbonyl; and
 R^3 and R^5 are hydrogen.


11. The thioacylating reagent according to claim 1, wherein the thioacylating reagent is selected from the group consisting of:

1-(α -Boc-L-alanyl-thioacyl)-2-benzimidazolone,
 1-(α -Boc-L-arginyl-di-N-Cbz-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-arginyl-N-tosyl-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-asparaginy-N-xanthenyl-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-aspartyl- β -benzyl ester-thioacyl)-2-benzimidazolone, 1-(α -Fmoc-L-aspartyl- β -t-butyl ester-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-cysteinyl-S-benzyl ether-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-glutamyl- γ -benzyl ester-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-glutaminy-N-xanthenyl-thioacyl)-2-benzimidazolone, 1-(α -Boc-glycyl-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-histidyl-N-


- 58 -

histidyl-N-tosyl-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-isoleucyl-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-leucyl-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-lysyl- ϵ -N-Cbz-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-methionyl-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-phenylalanyl-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-prolyl-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-threonyl-O-benzyl ether-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-tryphophyl-thioacyl)-2-benzimidazolone, 1-(α -Boc-L-tyrosinyl-O-2,6-dichlorobenzyl ether-thioacyl)-2-benzimidazolone, or 1-(α -Boc-L-valyl-thioacyl)-2-benzimidazolone.

12. An amino acid ortho amino thioanilide compound represented by the formula:




wherein R^1 is hydrogen or C_1 - C_4 branched or unbranched alkyl which may be substituted by a substituent selected from the group consisting of:



wherein A is -O- or -NH-,


D is benzyl or xanthenyl,

n is 1 or 2;




wherein E is -H or - CH_3 ,


- 59 -

G is $-\text{CH}_2-$, $-\text{O}-$ or $-\text{S}-$,J is $-\text{S}-$ or $-\text{CH}_2-$,L is $-\text{CH}_3$ or phenyl;

wherein M and T may be the same or different and are selected from the group consisting of hydrogen, fluorine, chlorine, bromine and iodine, and Q is hydrogen, hydroxy, or 2ClZ ,



wherein V is carbobenzyloxy or tosyl, or



R^2 is related from the group consisting of t-butoxy, carbobenzyloxy, chlorobenzyloxy, 9-fluorenylmethoxycarbonyl, tosyl, trityl and xanthenyl;

R^3 is hydrogen, methyl, or ethyl; or R^1 and R^3 , taken together with the carbon atom to which they are attached, form a cycloalkyl group containing 3-5 carbon atoms;

R^4 is hydrogen; or R^1 and R^4 , taken together with the carbon and nitrogen atoms to which they are

- 60 -


attached form a saturated heterocycle containing not less than 2, but not more than 6 ring carbon atoms; and

R^5 is selected from the class consisting of hydrogen, methyl, ethyl, flourine, chlorine, bromine, iodine, amino, amido, azido, hydroxy, hydroxymethyl, carboxy, carboxy methyl, cyano, guanido, mercapto, and nitro.

13. The amino acid ortho amino-thioanilide according to claim 12, wherein R^1 is C_1-C_4 branched or unbranched alkyl, R^2 is t-butoxy, and R^3 and R^5 are hydrogen.

14. The amino acid ortho amino thioanilide according to claim 12, wherein R^1 and R^4 , taken together with the carbon and nitrogen atoms to which they are attached form a saturated heterocycle containing 4 ring carbon atoms R^2 in the compound is t-butoxy, and R^3 and R^5 hydrogen.

15. The amino acid ortho amino thioanilide according to claim 12, wherein R^1 is represented by the formula:



wherein A is -O- or -NH-,

D is benzyl or xanthenyl,


n is 1 or 2;

R^2 is t-butoxy; and

R^3 and R^5 are hydrogen.

- 61 -

16. The amino acid ortho-amino-thioanilide according to claim 12, wherein R^1 is represented by the formula:



wherein E is -H or $-\text{CH}_3$,

G is $-\text{CH}_2-$, $-\text{O}-$ or $-\text{S}-$,


J is $-\text{S}-$ or $-\text{CH}_2-$,

L is $-\text{CH}_3$ or phenyl;

R_2 is t-butoxy; and

R_3 and R_5 are hydrogen.

17. The amino acid ortho amino thioanilide according to claim 12, wherein R_1 is represented by the formula:




wherein M and T may be the same or different and are members selected from the group consisting of hydrogen, fluorine, chlorine, bromine and iodine, and Q is hydrogen hydroxy, or 2Cl_2 ;

R^2 is t-butoxy; and

R^3 and R^5 are hydrogen.


18. The amino acid ortho amino thioanilide according to claim 12, wherein R^1 is represented by the formula:



- 62 -


wherein V is carbobenzyloxy or tosyl;
 R^2 is t-butoxy; and
 R^3 and R^5 are hydrogen.

19. The amino acid ortho amino thioanilide according to claim 12, wherein R^1 in the compound is:




R^2 is t-butoxy; and
 R^3 and R^5 are hydrogen.

20. The amino acid ortho amino thioanalide according to claim 12, wherein R^1 is represented by the formula:



wherein n is 1-4 and R^3 and R^5 are hydrogen.

21. A compound according to claim 12, wherein R^1 is:



R^2 is 9-fluorenylmethyloxycarbonyl; and
 R^3 and R^5 are hydrogen.


- 63 -

22. The amino acid ortho-amino-thioanalide according to claim 12, wherein the amino acid ortho amino thioanalide is selected from the group consisting of:


α -N-Boc-L-alanyl-ortho-amino-thioanilide, α -N-Boc-L-arginyl-di-N-Cbz-ortho-amino-thioanilide, α -N-Boc-L-arginyl-N-tosyl-ortho-amino-thioanilide, α -N-Boc-L-asparaginyl-N-xanthenyl-ortho-amino-thioanilide, α -N-Boc-aspartyl- β -benzyl ester-ortho-amino-thioanilide, α -N-Fmoc-aspartyl- β -t-butyl ester-ortho-amino-thioanilide, α -N-Boc-L-cysteinyl-S-benzyl ether-ortho-amino-thioanilide, α -N-Boc-L-glutamyl- γ -benzyl ester-ortho-amino thioanilide, α -N-Boc-L-glutamyl-N-xanthenyl-ortho-amino thioanilide, α -N-Boc-glycyl-ortho-amino thioanilide, α -N-Boc-L-histidyl-N-benzyl-ortho-amino thioanilide, α -N-Boc-L-histidyl-N-tosyl-ortho-amino thioanilide, α -N-Boc-L-isoleucyl-ortho-amino-thioanilide, α -N-Boc-L-leucyl-ortho-amino thioanilide, α -N-Boc- ϵ -N-2ClZ-L-lysyl-ortho-amino thioanilide, α -N-Boc-L-methionyl-ortho-amino-thioanilide, α -N-Boc-L-phenylalanyl-ortho-amino thioanilide, α -N-Boc-L-prolyl-ortho-amino-thioanilide, α -N-Boc-L-threonyl-O-benzyl ether-ortho-amino-thioanilide, α -N-Boc-L-tryptophyl-ortho-amino-thioanilide, α -N-Boc-L-tyrosinyl-O-2,6-dichlorobenzyl ether-ortho-amino-thioanilide, or α -N-Boc-L-valyl-ortho-amino-thioanilide.

- 64 -

23. An amino acid ortho-amino-anilide compound represented by the formula:



wherein R^1 is hydrogen or C_1 - C_4 branched or unbranched alkyl which may be substituted by a substituent selected from the group consisting of:



wherein A is -O- or -NH-,

D is benzyl or xanthenyl,

n is 1 or 2;




wherein E is -H or - CH_3 ,

G is - CH_2 -, -O- or -S-,


J is -S- or - CH_2 -,

L is - CH_3 or phenyl;




wherein M and T may be the same or different and are members selected from the group consisting of hydrogen, fluorine, chlorine, bromine and iodine, and Q is hydrogen, hydroxy, or $2ClZ$,

- 65 -



wherein V is carbobenzyloxy or tosyl, or



R² is selected from the group consisting of t-butoxy, carbobenzyloxy, chlorobenzyloxy, 9-fluorenylmethyloxycarbonyl, tosyl, trityl and xanthenyl;

R³ is hydrogen, methyl, or ethyl; or R¹ and R³, taken together with the carbon atom to which they are attached, form a cycloalkyl group containing 3-5 carbon atoms;

R⁴ is hydrogen; or R¹ and R⁴, taken together with the carbon and nitrogen atoms to which they are attached, form a saturated heterocycle containing not less than 2, but not more than 6 ring carbon atoms; and


R⁵ is selected from the group consisting of hydrogen, methyl, ethyl, fluorine, chlorine, bromine, iodine, amino, amido, azido, hydroxy, hydroxymethyl, carboxy, carboxymethyl, cyano, guanido, mercapto, and nitro.

24. The amino acid ortho amino anilide according to claim 23, wherein R¹ is C₁-C₄ branched or unbranched alkyl, R² is t-butoxy, and R³ and R⁵ are hydrogen.

- 66 -

25. The amino acid ortho-amino-anilide according to claim 23, wherein R^1 and R^4 taken together with the carbon and nitrogen atoms to which they are attached, form a saturated heterocycle containing 4 ring carbon atoms (a pyrrolidine ring), R^2 is t-butoxy, and R_3 and R_5 are hydrogen.

26. The amino acid ortho-amino-anilide according to claim 23, wherein R^1 is represented by the formula:



wherein A is -O- or -NH-,


D is benzyl or xanthenyl,

n is 1 or 2;

R^2 is t-butoxy; and

R^3 and R^5 are hydrogen.

27. The amino acid ortho amino anilide according to claim 23, wherein R^1 is represented by the formula:



wherein E is -H or -CH₃,

G is -CH₂-, -O- or -S-,

J is -S- or -CH₂-,


L is -CH₃ or phenyl;

R^2 is t-butoxy; and

R^3 and R^5 are hydrogen.

28. The amino acid ortho amino anilide according to claim 23, wherein R^1 is represented by the formula:

- 67 -




wherein M and T may be the same or different and are selected from group consisting of hydrogen, fluorine, chlorine, bromine and iodine, and Q is hydrogen, hydroxy, or 2ClZ;

R² is t-butoxy; and

R³ and R⁵ are hydrogen.

29. The amino acid ortho amino anilide according to claim 23, wherein R¹ is represented by the formula:




wherein V is carbobenzyloxy or tosyl;

R² is t-butoxy; and

R³ and R⁵ are hydrogen.

30. The amino acid ortho-amino-anilide according to claim 23, wherein R¹ is:




R² is t-butoxy; and

R³ and R⁵ are hydrogen.

;


- 68 -

31. The amino acid ortho-amino-anilide according to claim 23, wherein R^2 is represented by the formula:



wherein n is 1-4 and R^3 and R^5 are hydrogen.

32. The amino acid ortho-amino-anilide according to claim 23, wherein R^1 is:



R^2 is 9-fluorenylmethyloxycarbonyl; and
 R^3 and R^5 are hydrogen.


33. The amino acid ortho-amino-anilide according to claim 23, wherein the amino acid ortho-amino-anilide is selected from the group consisting of:

α -N-Boc-L-arginyl-di-N-Cbz-ortho-amino anilide,
 α -N-Boc-L-arginyl-N-tosyl-ortho-amino-anilide,
 α -N-Boc-L-asparaginyll-N-xanthenyl-ortho-amino-anilide,
 α -N-Boc-aspartyl- β -benzyl ester-ortho-amino-anilide,
 α -N-Fmoc-aspartyl- β -t-butyl ester-ortho-amino-anilide,
 α -N-Boc-L-cysteinyl-S-benzyl ether-ortho-amino-anilide,
 α -N-Boc-L-glutamyl- γ -benzyl ester-ortho-amino-anilide,
 α -N-Boc-L-glutaminyl-N-xanthenyl-ortho-amino-anilide,
 α -N-Boc-glycyl-ortho-amino-anilide, α -N-Boc-L-histidyl-N-benzyl-ortho-amino-anilide, α -N-Boc-L-histidyl-N-tosyl-ortho-amino-anilide, α -N-Boc-L-isoleucyl-ortho-amino-anilide, α -N-Boc-L-leucyl-ortho-amino-anilide, α -N-Boc- ϵ -N-2ClZ-L-lysyl-ortho-amino anilide, α -N-Boc-L-methionyl-ortho-amino-anilide, α -N-Boc-L-phenylalanyl-ortho-amino-anilide, α -N-Boc-L-prolyl-ortho-amino-anilide, α -N-Boc-L-threonyl-O-

- 69 -

benzyl ether-ortho-amino-anilide, α -N-Boc-L-tryptophyl-ortho-amino anilide, α -N-Boc-L-tyrosinyl-O-2,6-dichlorobenzyl ether-ortho-amino-anilide, or α -N-Boc-L-valyl-ortho-amino-anilide.

34. A peptide represented by the formula:



and salts thereof,

wherein R^1 , R^3 , and R^4 are defined as according to claim 1;

each X is, independently, oxygen or sulphur, provided that at least one atom represented by X is sulphur; and

n is 1-4.


35. The peptide according to claim 34, wherein one atom represented by X is sulphur and the remaining atoms represented by X are each, independently, oxygen.

36. The peptide according to claim 34, wherein n is 3.


37. The peptide according to claim 34, wherein n is 2.

- 70 -


38. The peptide according to claim 34,
having the formula:



39. The peptide according to claim 34,
having the formula:




40. The peptide according to claim 34,
having the formula:




- 71 -


41. The peptide according to claim 34,
having the formula:



42. The peptide according to claim 34,
having the formula:




43. The peptide according to claim 34,
having the formula:




- 72 -

44. The peptide according to claim 34, having the formula:




45. A process for selectively producing monosubstituted protected amino acid ortho-amino-anilides which comprises contacting ortho-phenylene diamine of the formula $C_6H_4(NH_2)_2$ with a protected amino acid or protected polypeptide in the presence of a peptide coupling agent in an inert organic solvent.

46. The process according to claim 45, wherein said amino acid ortho amino anilide is represented by the formula:



wherein R^1 is hydrogen or C_1 - C_4 branched or unbranched alkyl which may be substituted by a substituent selected from the group consisting of:





wherein A is -O- or -NH-,


D is benzyl or xanthenyl,

- 73 -


n is 1 or 2;

wherein E is -H or -CH₃,G is -CH₂-, -O- or -S-,J is -S- or -CH₂-,L is -CH₃ or phenyl;

wherein M and T may be the same or different and are selected from the group consisting of hydrogen, fluorine, chlorine, bromine and iodine, and Q is hydrogen, hydroxy, or 2ClZ,



wherein V is carbobenzyloxy or tosyl, or



R² is selected from the group consisting of t-butoxy, carbobenzyloxy, chlorobenzyloxy, 9-fluorenylmethoxycarbonyl, tosyl, trityl and xanthenyl;

- 74 -


R^3 is hydrogen methyl, or ethyl; or R^1 and R^3 , taken together with the carbon atom to which they are attached, form a cycloalkyl group containing 3-5 carbon atoms;

R^4 is hydrogen; or R^1 and R^4 , taken together with the carbon and nitrogen atoms to which they are attached, form a saturated heterocycle containing not less than 2, but not more than 6 ring carbon atoms; and


R^5 is selected from the group consisting of hydrogen, methyl, ethyl, fluorine, chlorine, bromine, iodine, amino, amido, azido, hydroxy, hydroxymethyl, carboxy, carboxymethyl, cyano, guanido, mercapto, and nitro.

47. The process according to claim 45, wherein said organic solvent is selected from the group consisting of dichloromethane, chloroform, N,N'-dimethylformamide and tetrahydrofuran.

48. The process according to claim 45, wherein said protected amino acid is represented by the formula:



wherein R^1 is hydrogen or C_1 - C_4 branched or unbranched alkyl which may be substituted by a substituent selected from the group consisting of:




wherein A is -O- or -NH-,

D is benzyl or xanthenyl,

n is 1 or 2;

- 75 -




wherein E is -H or -CH₃,


G is -CH₂-, -O- or -S-,

J is -S- or -CH₂-,


L is -CH₃ or phenyl;



wherein M and T may be the same or different and are selected from the group consisting of hydrogen, fluorine, chlorine, bromine and iodine, and Q is hydrogen, hydroxy, or 2ClZ,



wherein V is carbobenzyloxy or tosyl, or



R² is selected from the group consisting of t-butoxy, carbobenzyloxy, chlorobenzyloxy, 9-fluorenyl methyloxycarbonyl, tosyl, trityl, and xanthenyl;

R³ is hydrogen, methyl, or ethyl; or R¹ and R³, taken together with the carbon atom to which they are attached, form a cycloalkyl group containing 3-5 carbon atoms; and


- 76 -

R^4 is hydrogen; or R^1 and R^4 , taken together with the carbon and nitrogen atoms to which they are attached, form a saturated heterocycle containing not less than 2, but not more than 6 ring carbon atoms.


49. The process according to claim 45, wherein said peptide coupling agent is selected from the group consisting of N,N'-dicyclohexylcarbodiimide, N,N'-carbonyl diimidazole, 1-hydroxybenzotriazole, ethyl chloroformate and benzyl chloroformate.

50. A process for the selective production of amino acid ortho-amino-thioanilides which comprises contacting an amino acid ortho-amino-anilide with a reagent comprising a composition of phosphorous pentasulfide and sodium carbonate in an inert organic solvent.

51. The process according to claim 50, wherein said protected amino acid ortho-amino thioanilide is represented by the formula:



wherein R^1 is hydrogen or C_1 - C_4 branched or unbranched alkyl which may be substituted by a substituent selected from the group consisting of:





wherein A is -O- or -NH-,


D is benzyl or xanthenyl,

- 77 -


n is 1 or 2;

wherein E is -H or -CH₃,G is -CH₂-, -O- or -S-,J is -S- or -CH₂-,L is -CH₃ or phenyl;

wherein M and T may be the same or different and are selected from the group consisting of hydrogen, fluorine, chlorine, bromine and iodine, and Q is hydrogen, hydroxy, or 2ClZ,



wherein V is carbobenzyloxy or tosyl, or



R² is selected from the group consisting of t-butoxy, carbobenzyloxy, chlorobenzyloxy,

- 78 -

9-fluorenylmethyloxycarbonyl, tosyl, trityl and xanthenyl;


R^3 is hydrogen, methyl, or ethyl; or R^1 and R^3 , taken together with the carbon atom to which they are attached, form a cycloalkyl group containing 3-5 carbon atoms;

R^4 is hydrogen; or R^1 and R^4 , taken together with the carbon and nitrogen atoms to which they are attached, form a saturated heterocycle containing not less than 2, but not more than 6 ring carbon atoms; and

R^5 is selected from the class consisting of hydrogen, methyl, ethyl, fluorine, chlorine, bromine, iodine, amino, amido, azido, hydroxy, hydroxymethyl, carboxy, carboxymethyl, cyano, guanido, mercapto, and nitro.


52. The process according to claim 50, wherein said solvent is selected from the group consisting of diethyl ether and tetrahydrofuran.

53. The process for selective formation of protected 1-amino thioacylbenzimidizoyl-2-ones of the formula:




- 79 -


wherein R^1 is hydrogen or C_1 - C_4 branched or unbranched alkyl which may be substituted by a substituent selected from the group consisting of:




wherein A is -O- or -NH-,
D is benzyl or xanthenyl,
n is 1 or 2;



wherein E is -H or -CH₃,
G is -CH₂-, -O- or -S-,
J is -S- or -CH₂-,
L is -CH₃ or phenyl;




wherein M and T may be the same or different and are members selected from the group consisting of hydrogen, fluorine, chlorine, bromine and iodine, and Q is hydrogen, hydroxy, or 2ClZ,



wherein V is carbobenzyloxy or tosyl, or

- 80 -



R^2 is selected from the group consisting of t-butoxy, carbobenzyloxy, chlorobenzyloxy, 9-fluorenylmethyloxycarbonyl, tosyl, trityl and xanthenyl;


R^3 is hydrogen, methyl, or ethyl; or R^1 and R^3 , taken together with the carbon atom to which they are attached, form a cycloalkyl group containing 3-5 carbon atoms;

R^4 is hydrogen; or R^1 and R^4 , taken together with the carbon and nitrogen atoms to which they are attached, form a saturated heterocycle containing not less than 2, but not more than 6 ring carbon atoms; and


R^5 is selected from the group consisting of hydrogen, methyl, ethyl, fluorine, chlorine, bromine, iodine, amino, amido, azido, hydroxy, hydroxymethyl, carboxy, carboxymethyl, cyano, guanido, mercapto, and nitro which comprises contacting an amino acid ortho amino thioanilide, as defined according to claim 12, with carbonyl ditriazole in the presence of an inert etheral solvent.

- 81 -

54. A process for selective formation of protected 1-amino thioacylbenzimidazol-2-ones of the formula:




wherein R^1 is hydrogen or C_1 - C_4 branched or unbranched alkyl which may be substituted by a substituent selected from the group consisting of:



wherein A is -O- or -NH-,

D is benzyl or xanthenyl,

n is 1 or 2;




wherein E is -H or - CH_3 ,

G is - CH_2 -, -O- or -S-,


J is -S- or - CH_2 -,

L is - CH_3 or phenyl;


- 82 -



wherein M and T may be the same or different and are selected from the group consisting of hydrogen, fluorine, chlorine, bromine and iodine, and Q is hydrogen, hydroxy, 2ClZ,



wherein V is carbobenzyloxy or tosyl, or



R² is selected from the group consisting of t-butoxy, carbobenzyloxy, chlorobenzyloxy, 9-fluorenylmethyloxycarbonyl, tosyl, trityl and xanthenyl;

R³ is hydrogen, methyl, or ethyl; or R¹ and R³, taken together with the carbon atom to which they are attached, form a cycloalkyl group containing 3-5 carbon atoms;

R⁴ is hydrogen; or R¹ and R⁴, taken together with the carbon and nitrogen atoms to which they are attached, form a saturated heterocycle containing not less than 2, but not more than 6 ring carbon atoms; and

R⁵ is selected from the class consisting of hydrogen, methyl, ethyl, fluorine, chlorine, bromine, iodine, amino, amido, azido, hydroxy, hydroxymethyl,

- 83 -

carboxy, carboxymethyl, cyano, guanido, mercapto, and nitro

which comprises contacting on amino acid ortho amino thioanilide, as defined according to claim 12, with 2,4,6-collidine in the presence of an inert etheral solvent.

55. A process for selective formation of thioamide bonds which comprises contacting a protected amino acid or a protected peptide, said protected amino acid or protected peptide having a free amino functionality, with a thioacylating reagent, as defined according to claim 1, in the presence of an inert solvent.

56. A process for selective formation of thioamide bonds, wherein said process employs solid phase peptide synthesis techniques which comprises contacting a protected amino acid or protected peptide, said protected amino acid or protected peptide having a free amino terminus and suitably attached to an insoluable and inert resin by its carboxy terminus, with a thioacylating reagent, as defined according to claim 1, in the presence of an inert solvent.

57. The process according to claim 55 wherein said solvent is selected from the group consisting of acetonitrile, dichloromethane, diethyl ether, N,N'-dimethylformamide, dimethylsulfoxide and tetrahydrofuran.

58. The process according to claim 56, wherein said solvent is selected from the group consisting of acetonitrile, dichloromethane, diethyl

- 84 -

ether, N,N'-dimethylformamide, dimethylsulfoxide and tetrahydrofuran.


INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 90/00248

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ¹		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC ⁵ : C 07 D 235/26, C 07 K 5/08, C 07 K 5/10, C 07 K 5/06		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
IPC ⁵ : C 07 D 235/00, C 07 K 5/00, C 07 D 403/00, C 07 C 327/00		
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ¹⁰	Citation of Document, ¹¹ with Indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	Farmaco, Edizione Scientifica, volume 39, no. 4, 1984, (Pavia, IT), S. Salvadori et al.: "Opioid peptides. Structure-activity relationships of dermorphin endothiotetrapeptides", pages 316-321, see the whole article	34-36

X	Tetrahedron Letters, volume 28, no. 19, 1987, Pergamon Journals Ltd., (Oxford GB), David W. Brown et al.: "Mono- and dithionopeptide synthesis", pages 2171-2174, see the whole article	34-35

./.		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
30th October 1990	21 NOV. 1990	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	 MISS T. TAZELAAR	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, ¹⁾ with indication, where appropriate, of the relevant passages	Relevant to Claim No.
X	Chemical Abstracts, volume 102, no. 11, 18 March 1985, (Columbus, Ohio, US), G. Lajoie et al.: "Synthesis and biological activity of monothio- nated analogs of leucine-enkephalin", see page 615, abstract 96050q & Int. J. Pept. Protein Res. 1984, 24(4), 316-27 ---	34-35
X	Chemical Abstracts, volume 101, no. 7, 13 August 1984, (Columbus, Ohio, US), K. Clausen et al.: "Role of the peptide backbone in biological activity: synthesis of enkephalins with psi(CH ₂ S) and psi(CSNH) amide bond replacements", see page 660, abstract 55517h & Pept.: Struct. Funct., Proc. Am. Pept. Symp., 8th 1983, 307-10 ---	34-35
X	Chemical Abstracts, volume 100, no. 23, 4 June 1984, (Columbus, Ohio, US), K. Clausen et al.: "Evidence of a peptide backbone contribution toward selective receptor recognition for leucine enkephalin thioamide analogs", see page 75, abstract 185964s & Biochem. Biophys. Res. Commun. 1984, 120(1), 305-10 ---	34-35
X	Chemical Abstracts, volume 100, no. 9, 27 February 1984, (Columbus, Ohio, US), K. Clausen et al.: "Synthesis of leucine enkephalin and aspartame analogues containing thioamide linkages at specific positions", see page 638, abstract 68688u & Pept. Proc., Eur. Pept. Symp., 17th 1982 (Pub. 1983), 207-10 ---	34-35
X	Chemical Abstracts, volume 95, no. 23, 7 December 1981, (Columbus, Ohio, US), W.L. Mock et al.: "Hydrolysis of a thiopeptide by cadmium carboxypeptidase A.", see page 292, abstract 199699z & Biochem. Biophys. Res. Commun. 1981, 102(1), 389-96 ---	34-35
	./.	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, " with indication, where appropriate, of the relevant passages	Relevant to Claim No.
X	<p>Journal of the Chemical Society, Perkin Transactions I, (JCPRB4(4)), 1984 (Letchworth, GB), K. Clausen et al.: "Studies on amino acids and peptides. Part 6. Methods for introducing thioamide bonds into the peptide backbone: Synthesis of the four monothio analogues of leucine enkephalin", pages 785-798 see the whole article cited in the application</p> <p>---</p>	34-35
X	<p>Journal of the American Chemical Society, volume 104, no. 19, 22 September 1982, American Chemical Society, (Gaston, PA, US), P. Campbella et al.: "Carboxypeptidase A catalyzed hydrolysis of thiopeptide and thionester analogues of specific substrates. An effect on Kcat for peptide, but not ester, substrates", pages 5221-5226 see the whole article cited in the application</p> <p>---</p>	34-35
A	<p>Journal of Polymer Science: Part A: Polymer Chemistry, Polymer Chemistry Edition, volume 27, no. 5, April 1989, John Wiley and Sons, (New York, US), A.R. Katritzky et al.: " Azlactones as polymer components and intermediates", pages 1781-1790 see the whole article</p> <p>---</p>	
A	<p>EP, A, 0222283 (MERCK PATENT GESELLSCHAFT MIT BESCHRANKTER HAFTUNG) 20 May 1987</p> <p>---</p>	
P,X	<p>Journal of the American Chemical Society, volume 112, no. 1, 3 January 1990, American Chemical Society, (Gaston, PA, US), D. B. Sherman et al.: " Compatibility of thioamides with reverse turn features: Synthesis and conformational analysis of two model cyclic pseudo-peptides containing thioamides as backbone modifications", pages 433-441 see the whole article</p> <p>-----</p>	34-35

CA 9000248
SA 38916

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0222283	20-05-87	DE-A- 3540495	21-05-87
		AU-B- 588929	28-09-89
		AU-A- 6462986	21-05-87
		JP-A- 62187488	15-08-87
		US-A- 4721776	26-01-88
<hr/>			

F.I.C. F()RM 10479

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.